arXiv:1210.7221v1 [math.OC] 26 Oct 2012

The value of Markov Chain Games with incomplete information on both sides

Fabien Gensbittel[*], Jérôme Renault[*]

October 29, 2012

Abstract

We consider zero-sum repeated games with incomplete information on both sides, where the states privately observed by each player follow independent Markov chains. It generalizes the model, introduced by Aumann and Maschler in the sixties and solved by Mertens and Zamir in the seventies, where the private states of the players were fixed. It also includes the model introduced in Renault [20], of Markov chain repeated games with lack of information on one side, where only one player privately observes the sequence of states. We prove here that the limit value exists, and we obtain a characterization via the Mertens-Zamir system, where the "non revealing value function" plugged in the system is now defined as the limit value of an auxiliary "non revealing" dynamic game. This non revealing game is defined by restricting the players not to reveal any information on the *limit behavior* of their own Markov chain, as in Renault 2006. There are two key technical difficulties in the proof: 1) proving regularity, in the sense of equicontinuity, of the T-stage non revealing value functions, and 2) constructing strategies by blocks in order to link the values of the non revealing games with the original values.

Key words: repeated games; incomplete information; zero-sum games; Markov chain; lack of information on both sides; stochastic games; Mertens-Zamir system.

1 Introduction

The pioneering work of Aumann and Maschler in the 1960s introducing Repeated games with incomplete information (see [1] for a re-edition of their work) was widely studied and extended. One of their most famous results concerns the model of two-player zero-sum repeated game with lack of information on one side and perfect observation: At the beginning of the game, a state variable k is chosen at random in a finite set K using some probability p, and announced to Player 1 only. A finite zero-sum game G_k depending on the state variable is then played repeatedly and after each stage the actions played are observed by both players. The famous

---

[*] TSE (GREMAQ, Université Toulouse 1 Capitole), 21 allée de Brienne, 31000 Toulouse, France. E-mails: fabien.gensbittel@tse-fr.eu, jerome.renault@tse-fr.eu

cav$u$ theorem of Aumann and Maschler then reads as follows: the value of the infinitely repeated game exists and is characterized as the concave hull of the value of an auxiliary game called the non-revealing game. In a previous work of Renault [20], this model was extended in the following direction: instead of being fixed once for all at the beginning of the game, the state variable was assumed to evolve according to a Markov chain of initial probability p and transition matrix M over K. Renault proved that in this model, called Markov chain game with incomplete information on one side, the value of the infinitely repeated game exists and gave a characterization for it, very close to the one of Aumann and Maschler. The main difference was in the definition of the non-revealing game: it was defined in Renault [20] as a dynamic game with infinite length where Player 1 was constrained not to reveal any information on the asymptotic behavior of the Markov chain rather than on the current state variable itself. The model of Markov chain game with incomplete information on one side was also studied by Neyman [16] who proved the existence of an optimal strategy for Player 1 and also proved a generalization of the existence of the value in case of imperfect observation of actions.

A more involved model also introduced in [1] is the model of two-player zero-sum repeated game with lack of information on both sides and perfect observation: At the beginning of the game, two states of nature k and l are chosen independently in some finite sets K and L, according to some probabilities p and q. The state k is announced to Player 1 only, and the state l is announced to Player 2 only. These states of nature determine a finite zero-sum game $G_{k,l}$ which is then played repeatedly; after each stage the actions played are observed by both players. Aumann and Maschler showed that the infinitely repeated game may have no value. Mertens and Zamir [14], showed the existence of the limit value (i.e. the existence of a limit for the sequence of values of finitely repeated games when the number of repetitions goes to infinity) in this model and gave a characterization for it (see the system (1) later) based on a system of functional equations acting on the value of the non-revealing game.

The results of Aumann and Maschler and of Mertens and Zamir led to a great number of works dedicated to generalizations of this model, or close extensions of it. Let us cite for games with incomplete information on one side, Kohlberg [10] for an explicit construction of an optimal strategy for Player 2, De Meyer and Rosenberg [4] and Laraki [12] for alternative proofs of existence of the limit value based on duality, Gensbittel [5] for the extension to infinite action spaces, or Sorin [26], Hart [7], Simon *et al.* [23], Renault [18] [19] for the nonzero-sum case.

Concerning incomplete information on both sides, let us mention the proof of existence of asymptotically optimal strategies by Heuer [8], the extension of the Mertens-Zamir system by Sorin [27], the study of an abstract game called the splitting game related to the system of functional equations by Laraki [13],alternative proofs of existence of the limit value based on the so-called operator approach by Rosenberg and Sorin [22], or more recently on continuous-time approach by Cardaliaguet, Laraki and Sorin [3].

We only consider here two-player zero-sum games and generalize the model of repeated game with lack of information on both sides to the case where the states of nature (k, l) are no longer fixed at the beginning of the game, but evolve according to given independent Markov chains $(k_t)_{t\geq 1}$ and $(l_t)_{t\geq 1}$. At the beginning of each stage, k_t is observed by Player 1 only, and l_t is observed by Player 2 only. We call such games Markov chain games with lack of information on both sides. Note that this model admits as a special case the model of repeated game with incomplete information on both sides, in which the value of the infinitely repeated game may not exist. In this paper, we generalize both the proofs of Renault [20] and Mertens and Zamir [14] and show the existence of the limit value for Markov chain games with lack of information on both sides. We also give a characterization for this limit value based on a system of functional equations, similar to the one introduced by Mertens and Zamir, and on the generalized notion of non-revealing games as introduced by Renault.

However, as already mentioned in these two works, our expression of the value cannot be easily computed from the basic data of the game. It was already noticed in [14], where several examples of computations are given, underlying that the limit value may be outside of the class of semi-algebraic functions, which contrasts with properties of the limit value for stochastic games (with complete information and finite state and action spaces). Moreover, the problem of computation of the value of non-revealing games for Markov chain games with incomplete information on one side was already mentioned by Renault who gave an example which appeared to be very difficult to compute, except for very particular values of the transition matrix as shown in the work of Hörner *et al.* [9].

Section 2 of this paper contains the model. Section 3 contains preliminary results and notations. A few important examples, and the main ideas of the proof, are presented in section 4. In section 5, we define and study a notion of nonrevealing strategies via projection matrices that will be used to quantify a notion of relevant information for both players. We introduce the auxiliary games called nonrevealing games, where both players are restricted to play a nonrevealing strategy, and prove that they have a limit value. In section 6, we introduce a system of functional equations very similar to the system introduced by Mertens and Zamir and prove that the limit value is the unique solution of this system, which is itself parameterized by the limit value of the nonrevealing game. Section 7 contains the proof of some technical results. We conclude in section 8 with a list of open questions related to the possible extensions of the model.

2 Model

Given a finite set S, $\Delta(S)$ denotes the set of probabilities over S and the cardinal of S is, with a slight abuse of notations, also denoted by S. The set of positive integers is denoted $\mathbb{N}^*$.

We consider a zero-sum game between 2 players with action sets I and J, sets of states K and L, where I, J, K, L are disjoint finite non empty sets, and payoff function

$$g : K \times L \times I \times J \longrightarrow [-1, 1].$$

M and N are given Markov matrices on K and L respectively, i.e. $M = (M_{k,k'})_{k,k' \in K}$ is a $K \times K$ matrix with non-negative entries and such that for all $k \in K$, $\sum_{k' \in K} M_{k,k'} = 1$, and similarly for N. The letters $p \in \Delta(K)$ and $q \in \Delta(L)$ denote initial probabilities.

The game is played by stages in discrete time. We are given two independent Markov chains: $(k_t)_{t \geq 1}$ with initial distribution p and transition matrix M, and $(l_t)_{t \geq 1}$ with initial distribution q and transition matrix N. At the beginning of every stage $t \geq 1$, Player 1 observes k_t and Player 2 observes l_t. Then both players simultaneously select an action in their action set, if (i_t, j_t) in $I \times J$ is played then Player 1's payoff for stage t is $g(k_t, l_t, i_t, j_t)$. Then (i_t, j_t) is publicly observed and the play goes to stage $t + 1$. Notice that the payoff

$g(k_t, l_t, i_t, j_t)$ is not directly observed and may not be deduced by the players at the end of stage t.

We denote by Σ and $\mathcal{T}$ the sets of behavior strategies of the players. Formally, for $t \geq 1$, let $H_t = (I \times J)^t$ denote the set of histories of actions of length t, with the convention $H_0 = \{\emptyset\}$. A strategy $\sigma \in \Sigma$ is a sequence $(\sigma_t)_{t\geq 1}$, where σ_t is a map from $H_{t-1} \times K^t$ to $\Delta(I)$. For $T \in I\!N^*$, the set Σ_T denotes the set of T-stage strategies for Player 1, that is of finite sequences $(\sigma_1, .., \sigma_T)$ induced by elements of Σ. $\mathcal{T}$ and $\mathcal{T}_T$ are defined similarly.

The value of the T-stage game is denoted $v_T(p, q)$, i.e.

$$v_T(p, q) = \max_{\sigma\in\Sigma} \min_{\tau\in\mathcal{T}} \gamma_T^{p,q}(\sigma, \tau) = \min_{\tau\in\mathcal{T}} \max_{\sigma\in\Sigma} \gamma_T^{p,q}(\sigma, \tau),$$

where $\gamma_T^{p,q}(\sigma, \tau) = I\!E_{p,q,\sigma,\tau} \frac{1}{T} \sum_{t=1}^{T} g(k_t, l_t, i_t, j_t)$, the expectation being taken with respect to the probability $I\!P_{p,q,\sigma,\tau}$ induced by (p, q, σ, τ) on the set of plays $\Omega = (K \times L \times I \times J)^\infty$. Elements of $\Delta(K)$ are seen as row vectors, so that if p' in $\Delta(K)$ is the law of the state k_T of stage T, then $p'M$ is the law of the following state k_{T+1} (similarly elements of $\Delta(L)$ are seen as row vectors). We have the standard recursive formula, for all $T \geq 0$:

$$\begin{aligned}(T+1)v_{T+1}(p, q) &= \max_{x\in\Delta(I)^K} \min_{y\in\Delta(J)^L} \left(G(p, q, x, y) + T \sum_{i\in I, j\in J} x(p)(i)y(q)(j)v_T(\vec{p}(x, i)M, \vec{q}(y, j)N) \right) \\ &= \min_{y\in\Delta(J)^L} \max_{x\in\Delta(I)^K} \left(G(p, q, x, y) + T \sum_{i\in I, j\in J} x(p)(i)y(q)(j)v_T(\vec{p}(x, i)M, \vec{q}(y, j)N) \right).\end{aligned}$$

where $G(p, q, x, y) = \sum_{k,l,i,j} p^k q^l x^k(i) y^l(j) g(k, l, i, j)$,

$$\forall i \in I,\ x(p)(i) = \sum_k p^k x^k(i),\ \text{and}\ \vec{p}(x, i) = \left(\frac{p^k x^k(i)}{x(p)(i)}\right)_{k\in K} \in \Delta(K)\ \text{if}\ x(p)(i) > 0,$$

$$\forall j \in J,\ y(q)(j) = \sum_l q^l y^l(j),\ \text{and}\ \vec{q}(y, j) = \left(\frac{q^l y^l(j)}{y(q)(j)}\right)_{l\in L} \in \Delta(L)\ \text{if}\ y(q)(j) > 0,$$

and $\vec{p}(x, i)$ (resp. $\vec{q}(y, j)$) is defined arbitrarily if $x(p)(i) = 0$ (resp. $y(q)(j) = 0$). If at some stage t the belief of Player 2 on the current state k_t in K is represented by p, and if Player 1 plays the mixed action x^k if his current state is k, then $x(p)(i)$ will be the probability that a given action i in I is played at this stage. After observing i, $\vec{p}(x, i)$ represents the new belief of Player 2 on k_t, and $\vec{p}(x, i)M$ is the belief of Player 2 on k_{t+1}.

Theorem 2.1. *$v(p, q) = \lim v_T(p, q)$ exists.*

Moreover, we will provide a characterization for the limit function v.

Definition 2.2. *Let f be a function from $\Delta(K) \times \Delta(L)$ to $I\!R$.*

- *f is I-concave if for all $q \in \Delta(L)$ the function $p \mapsto f(p, q)$ is concave on $\Delta(K)$,*
- *f is II-convex if for all $p \in \Delta(K)$, the function $q \mapsto f(p, q)$ is convex on $\Delta(L)$,*
- *f is balanced if for all $(p, q) \in \Delta(K) \times \Delta(L)$:*

$$w(p, q) = w(pM, qN).$$

Notation 2.3. *We denote by* $\mathcal{W}$ *the set of continuous functions* $w : \Delta(K) \times \Delta(L) \longrightarrow [-1, 1]$.

We will soon assume (w.l.o.g., see sections 3 and 4) that both Markov chains on K and L are recurrent and aperiodic. In this case we will show in Theorem 6.1 that v can be expressed as:

$$\begin{aligned} v &= \inf \left\{ \begin{matrix} w \in \mathcal{W}, w \text{ is } I\text{-concave, balanced and} \\ \forall (p,q) \in \Delta(K) \times \Delta(L),\ w(p,q) \geq \text{vex}_{II}\text{Max}(w, \hat{v})(p,q) \end{matrix} \right\}, \\ &= \sup \left\{ \begin{matrix} w \in \mathcal{W}, w \text{ is } II\text{-convex, balanced and} \\ \forall (p,q) \in \Delta(K) \times \Delta(L),\ w(p,q) \leq \text{cav}_{I}\text{Min}(w, \hat{v})(p,q) \end{matrix} \right\}, \end{aligned}$$

where $\hat{v}$ is the “nonrevaling limit value” that will be defined in Proposition 5.17 and where cav_I and vex_{II} denote respectively the concave and convex hull with respect to the first and second variable (see Notation 3.3 below).

3 Preliminaries

In all the paper the sets $\Delta(K)$, $\Delta(L)$ and $\Delta(K) \times \Delta(L)$ are endowed with the L^1-norm, $\|p - p'\| = \sum_k |p^k - p'^k|$, $\|(p,q) - (p',q')\| = \sum_k |p^k - p'^k| + \sum_l |q^l - q'^l|$ etc...

Lemma 3.1. *For each* T, v_T *is* I*-concave,* II*-convex and 1-Lipschitz.*

Proof. The game $\Gamma_T(p,q)$ can be seen as a game with incomplete information on one side (the state space being either K or L). The result follows therefore from Propositions 2.1 and 2.2 in [25]. □

Lemma 3.2. *For all* $T \geq 1$*, and* $(p,q) \in \Delta(K) \times \Delta(L)$*, we have:*

$$|v_{T+1}(p,q) - v_T(p,q)| \leq \frac{2}{T}, \ \ |v_{T+1}(p,q) - v_T(pM, qN)| \leq \frac{2}{T},$$

and consequently $|v_T(p,q) - v_T(pM,qN)| \leq \frac{4}{T}$.

Proof. Let $\sigma \in \Sigma$ be an optimal strategy in the game $\Gamma_T(p,q)$, then for all $\tau \in \mathcal{T}$,

$$\gamma^{p,q}_{T+1}(\sigma, \tau) \geq \frac{T}{T+1} v_T(p,q) - \frac{1}{T+1} \geq v_T(p,q) - \frac{2}{T},$$

which implies $v_{T+1}(p,q) \geq v_T(p,q) - \frac{2}{T}$. The reverse inequality is obtained by inverting the role of the players. Using now the recursive formula, let x be defined by $x^k(i) = \frac{1}{I}$ for all $(k,i) \in K \times I$. By construction $\vec{p}(x,i) = p$ for all $i \in I$ and we have

$$\begin{aligned} v_{T+1}(p,q) &\geq -\frac{1}{T+1} + \frac{T}{T+1} \inf_{y \in \Delta(J)^K} \sum_{j \in J} y(q)(j) v_T(pM, \vec{q}(y,j)N) \\ &\geq -\frac{1}{T+1} + \frac{T}{T+1} v_T(pM, qN), \end{aligned}$$

where the last inequality follows from Jensen's inequality since v_T is II-convex and by construction $\sum_{j \in J} y(q)(j) \vec{q}(y,j) = q$. The reverse inequality is obtained by inverting the role of the players. The last statement follows from the triangle inequality. □

If some state k is transient, then for each ε there exists T_0 such that $I\!P(\forall T \geq T_0,\ k_T \neq k) \geq 1-\varepsilon$. Both players can wait until stage T_0 and enter a game where state k has disappeared with high probability, so we can assume w.l.o.g. that there is no transient state. From now on, we assume that all states in K and L are recurrent.

We now take care of periodicity. Some states may *a priori* be periodic, let T_0 be a common multiple of all periods in the chains induced by M and N. From the preceding lemma, we have $\limsup_T v_{TT_0}(p,q) = \limsup_T v_T(p,q)$, and $\liminf_T v_{TT_0}(p,q) = \liminf_T v_T(p,q)$, so to study $\lim_T v_T$ we can consider plays by blocks of T_0 stages (so that one stage now corresponds to T_0 original stages). The Markov chains with transitions matrices M^{T_0} and N^{T_0} are aperiodic, and this consideration is without loss of generality.

Summing up, from now on[1] we assume w.l.o.g. that both Markov chains are recurrent and aperiodic. We still use the letters M and N to denote the (now recurrent aperiodic) transitions matrices. We have the convergence of M^t to a stochastic matrix B such that $B^2 = B = BM = MB$. The state space K can be partitioned into recurrence classes $K(1)$, ..., $K(r_M)$, where r_M is the number of recurrence classes associated to M. Each class $K(r)$ has a unique invariant measure $p^*(r)$, and the k-th row of B corresponds to $p^*(r)$ where $K(r)$ is the class containing k. Invariant measures for B and M coincide, the set of those measures is the convex hull of $\{p^*(1), ..., p^*(r_M)\}$ denoted by P^*. Although B is not invertible if there are less recurrence classes than states, for p^* in P^* we write $B^{-1}(p^*) = \{p \in \Delta(K), pB = p^*\}$.

Regarding the Markov chain associated to the states in L, we proceed similarly and use the following notations. There is convergence of N^t to a stochastic matrix C. The state space L is partitioned into recurrence classes $L(1)$, ..., $L(r_L)$. Each class $L(r)$ has a unique invariant measure $q^*(r)$, and the l-th row of C corresponds to $q^*(r)$ where $L(r)$ is the class containing l. Invariant measures for C and N coincide, the set of those measures is the convex hull of $\{q^*(1), ..., q^*(r_L)\}$ denoted by Q^*. For q^* in Q^*, we write $C^{-1}(q^*) = \{q \in \Delta(K), qC = q^*\}$.

We will extensively use the following notations.

Notation 3.3. *If $f : X \longrightarrow I\!R$ is a bounded function defined on some convex set X, $\mathrm{cav}(f)$ denotes the smallest concave function greater or equal to f, and $\mathrm{vex}(f)$ denotes the largest convex function lower or equal to f. If f is defined on a product of convex sets $X \times Y$, for all $(x,y) \in X \times Y$, $\mathrm{cav}_I(f)(x,y)$ denotes the concave hull of the partial function $f(.,y)$ at point x. $\mathrm{vex}_{II}(f)$ is defined symmetrically.*

Notation 3.4. *For any continuous function $f : \Delta(K) \times \Delta(L) \longrightarrow I\!R$, we write:*

$$\mathcal{W}^+(f) = \left\{ \begin{array}{c} w \in \mathcal{W}, w \text{ is } I\text{-concave and} \\ \forall (p,q) \in \Delta(K) \times \Delta(L),\ w(p,q) \geq \mathrm{vex}_{II}\mathrm{Max}(w,f)(p,q) \end{array} \right\}$$

$$\mathcal{W}^-(f) = \left\{ \begin{array}{c} w \in \mathcal{W}, w \text{ is } II\text{-convex and} \\ \forall (p,q) \in \Delta(K) \times \Delta(L),\ w(p,q) \leq \mathrm{cav}_I\mathrm{Min}(w,f)(p,q) \end{array} \right\},$$

4 Examples and overview of the proof

Let us at first emphasize that, in contrast with repeated games with incomplete information, if Player 1 does not use his information on the states $(k_t)_{t\geq 1}$, beliefs of Player 2 over states evolve across stages according to the transition matrix M. Precisely, under this assumption, the belief of Player 2 on the current state k_{t+1} at stage $t+1$ (i.e., the conditional law of k_{t+1} given the actions played up to stage t) is simply the law of k_{t+1}, which is pM^t. Symmetrically, if Player 2 does not use his information on the states $(l_t)_{t\geq 1}$, then the belief of Player 1 on l_{t+1} at stage $t+1$ is qN^t.

---

[1] except in the examples D and E of section 4.

Example A: A Markov chain without memory. If both Markov chains are i.i.d. sequences, then the problem reduces to a classical repeated game, by replacing the payoff function in pure strategies by its expectation with respect to (k_t, l_t). The value of the infinitely repeated game exists and equals the value of this auxiliary one-stage game.

Example B: Irreducible aperiodic Markov chains. Assume that both chains are irreducible and aperiodic, say

$$K = L = \{a, b\}, \quad M = \begin{pmatrix} \frac{2}{3} & \frac{1}{3} \\ \frac{1}{3} & \frac{2}{3} \end{pmatrix}, N = \begin{pmatrix} \frac{3}{4} & \frac{1}{4} \\ 1 & 0 \end{pmatrix}.$$

This Markov chain has a unique recurrence class, so $r_M = r_N = 1$ and $K = K(1) = L = L(1)$. Unique invariant probabilities are $p^* = (\frac{1}{2}, \frac{1}{2})$ and $q^* = (\frac{4}{5}, \frac{1}{5})$. M^t and N^t respectively converge to:

$$B = \begin{pmatrix} \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & \frac{1}{2} \end{pmatrix} \text{ and } C = \begin{pmatrix} \frac{4}{5} & \frac{1}{5} \\ \frac{4}{5} & \frac{1}{5} \end{pmatrix}.$$

In this case, the limit value exists, and we even have the stronger result that the (uniform) value of the infinitely repeated game exists. At first, if Player 1 plays independently of the states, the beliefs of Player 2 converge to p^*. On the other hand, after any sufficiently long number of stages, he can always forget the past actions of Player 2 and act as if his beliefs were very close to q^*. Choose now some integer T_0 such that M^{T_0} and N^{T_0} are respectively close to B and C, and then T_1 much larger than T_0. Let us construct a strategy of Player 1 as follows: (1) play any strategy independent of the states during T_0 stages, then (2) play an optimal strategy in the game $\Gamma_{T_1}(p^*, q^*)$, and (3) come back to (1). Such a strategy guarantees to Player 1 $v_{T_1}(p^*, q^*)$ up to some arbitrarily small error, hence by choosing T_1 large enough Player 1 guarantees $\text{limsup}_T v_T(p^*, q^*)$ in the infinitely repeated game. Inverting the role of the players implies that the value of the infinitely repeated game exists and equals $\lim_T v_T(p^*, q^*)$. In particular it does not depend on the initial probabilities (p, q). Notice that this proof can be generalized to the case of any irreducible aperiodic Markov chain.

Example C: M and L are the identity matrix. In this case, the problem is a repeated game with incomplete information on both sides as studied by Mertens and Zamir [14] since the initial state is selected at the beginning of the game and remains constant. In terms of Markov chains, the states can be identified with the recurrence classes, and therefore any information on the states is an information on the asymptotic behavior of the corresponding chain.

Let us at first define the non-revealing game: it denotes the game where none of the players is allowed to use his information on the state variable (i.e. players are restricted tu use strategies which do not depend on the selected states). For each pair (p, q) of initial probabilities, this game can be analyzed as the repetition of the average matrix game $\sum_{k,l} p^k q^l g(k, l, ., .)$.

Consequently, its value is just the value $u(p,q) = \max_{x\in\Delta(I)} \min_{y\in\Delta(J)} \sum_{k,l} p^k q^l g(k,l,x(i),y(j))$ of the one-stage average game.

Mertens and Zamir proved that the limit value v is known to exist and is characterized as the unique function $w \in \mathcal{W}$ solution of the so-called Mertens-Zamir system:

$$\forall (p,q) \in \Delta(K)\times\Delta(L), \begin{cases} w(p,q) = \text{vex}_{II}\text{Max}(w,u)(p,q) \\ w(p,q) = \text{cav}_I \text{Min}(w,u)(p,q) \end{cases}. \tag{1}$$

This system can be recast in a system of functional inequalities: Mertens and Zamir also proved that

$$v = \sup\{w | w \in \mathcal{W}^-(u)\} = \inf\{w | w \in \mathcal{W}^+(u)\}.$$

The proof of the second equality was based [14] on game considerations, and as a consequence they obtained that $\sup\{w|w \in \mathcal{W}^-(f)\} = \inf\{w|w \in \mathcal{W}^+(f)\}$ for any continuous function f. This result was later established and extended by the same authors [15], using analytical arguments.

Assuming the equality $\sup\{w|w \in \mathcal{W}^-(u)\} = \inf\{w|w \in \mathcal{W}^+(u)\}$, the proof of convergence of (v_T) by Mertens and Zamir can be sketched as follows.

The main point is to show that for any function w in $\mathcal{W}^-(u)$, Player 1 can defend in the game of length n with initial probabilities p and q, the quantity $w(p,q)$ up to an error going to zero with N uniformly with respect to (p,q) and the chosen strategy of Player 2. This can be done as follows. The strategy of Player 2 being known, Player 1 can compute after each stage t the current pair of beliefs (p_t,q_t) of both players: p_t denotes the conditional law of k_{t+1} given past actions and q_t denotes the conditional law of l_{t+1} given past actions. Then, either $u(p_t,q_t) \geq w(p_t,q_t)$ and he plays an optimal strategy in the non-revealing game at (p_t,q_t), or $u(p_t,q_t) < w(p_t,q_t)$ and he can split his information (think of Player 1 sending a, possibly random, message to Player 2) in order to drive Player 2's beliefs at some points $\tilde{p}_t$ such that $u(\tilde{p}_t,q_t) \geq w(\tilde{p}_t,q_t)$. The existence of such a splitting is ensured by: 1) the property $w \in \mathcal{W}^-(u)$ and 2) the splitting Lemma (see e.g. the Lemmas 6.5 and 6.6 in the present work). This strategy would defend $w(p_t,q_t)$ at stage $t+1$ if Player 2 was not using his information on the state at stage $t+1$. This is almost the case since the error can be bounded by the expected L_1-variation of the beliefs $\|q_{t+1} - q_t\|$. Recall that the sequence of beliefs (q_t) is a martingale and a classical bound on its L_1 variation allows to bound uniformly the average error by a quantity vanishing with n. Moreover using the properties of w and the construction of the strategy of Player 1, the expected value of $w(p_t,q_t)$ is always greater or equal than $w(p,q)$. All this implies that $\text{liminf}_n v_n(p,q) \geq w(p,q)$, so we obtain $\text{liminf}_n v_n \geq \sup\{w|w \in \mathcal{W}^-(u)\}$. Inverting the role of the players, we also have $\text{limsup}_n v_n \leq \inf\{w|w \in \mathcal{W}^+(u)\}$ which concludes the proof.

Example D: A periodic chain Let K be $\{a,b\}$, and M be $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$.

This Markov chain has a unique recurrence class, which is periodic with period 2. The relevant information for Player 2 is not the recurrence class, but the initial state which determines whether the sequence of states $(k_t)_{t\geq 1}$ will be $\{a,b,a,b,...\}$ or $\{b,a,b,a,...\}$. By considering the auxiliary game in which stages are blocks of length 2 of the initial game, then the new transition matrix is M^2, hence the identity, and each initial state becomes now a recurrence class. The problem can be reduced to that of Example C. As explained in section

3, this idea of playing stages by blocks of fixed length can be used in the general case to assume without loss of generality that the chain is aperiodic.

Existence of transient states does not play an important role here. We, however, present now an example with a transient state which will be used later (see Remark 6.3).

Example E: $K = \{a, b, c\}$, and

$$M = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ \frac{1}{2} & \frac{1}{4} & \frac{1}{4} \end{pmatrix}.$$

The matrix $B = \lim_t M^t$ would in this case be the following:

$$B = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ \frac{2}{3} & \frac{1}{3} & 0 \end{pmatrix}.$$

The last example below will be used to illustrate several definitions.

Example F: $K = L = \{a, b, c\}$, and

$$M = N = \begin{pmatrix} \frac{2}{3} & \frac{1}{3} & 0 \\ \frac{1}{3} & \frac{2}{3} & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

In this case, there are two recurrence classes $K(1) = \{a, b\}$ and $K(2) = \{c\}$. The associated invariant measures are $p^*(1) = (\frac{1}{2}, \frac{1}{2}, 0)$ and $p^*(2) = (0, 0, 1)$, and the limit projection matrix is

$$B = \begin{pmatrix} \frac{1}{2} & \frac{1}{2} & 0 \\ \frac{1}{2} & \frac{1}{2} & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

The following figure illustrates some of our notations.



Existence of the limit value in case of the last Example F appears to be more difficult, and we do not know how to proceed without following the general proof presented in this paper. Following the general scheme of Renault [20], we consider two kinds of information: long-term information (corresponding to the recurrence classes of the two chains, as in the Mertens-Zamir case), and short-term information (corresponding to the state variable itself within some recurrence class).

We now come back to the general case, assuming the Markov chains are recurrent and aperiodic, and describe the different steps of our proof.

Overview of the Proof

By analogy with the proof in Renault [20], we define a non-revealing game, where Players are restricted to strategies that do not reveal information on the recurrence classes of the chains. Precisely, the strategy of Player 1 has to be such that, after each history of actions of length t, the conditional probability that k_{t+1} belongs to any recurrence class $K(r)$ remains unchanged (see Definition 5.3).

This conceptual definition has to be compared with the non-revealing strategies introduced by Aumann and Maschler in case each player observes after each stage a signal which is a function of the actions played, and which requires that the beliefs of Player 2 on the state k remains unchanged.

The main difference here is that the study of the non-revealing game can not be reduced to that of a finite matrix game. Following the scheme of the proof of example B, with appropriate modifications, we show that the limit value $\hat{v}$ of this non-revealing game exists and that it depends only on the limit distributions of the two Markov chains (pB, qC). This proof requires however a precise analysis of the T-stage non-revealing games, for which we prove the existence of the value $\hat{v}_T$ and an appropriate recursive formula. The crucial point of the analysis of the non revealing values is to establish that the family $(\hat{v}_T)_{T\geq 1}$ is uniformly equi-continuous (this is the aim of the whole section 7), which implies uniform convergence of these values to their limit.

The second main technical difficulty is to link the value of the non revealing games with those of the original game, i.e. to adapt the proof of Mertens and Zamir illustrated in example C. We use the uniform convergence mentioned above, in particular for sufficiently large T_0, $\hat{v}_{T_0}$ is ε-close to $\hat{v}$. We prove that for any $N \in I\!N^*$, and for any *balanced* function $w \in \mathcal{W}^-(\hat{v})$, Player 1 can defend $w(p,q)$ in $\Gamma_{NT_0}(p,q)$ up to some error going to zero with n uniformly with respect to (p,q) and the chosen strategy of Player 2. The construction of the strategy of Player 1 is the same as explained in example C, except that stages are replaced by blocks of T_0 stages: Player 1 can compute after each block the current pair of beliefs (p_{nT_0}, q_{nT_0}) of both players and either (1) $\hat{v}(p_{nT_0}, q_{nT_0}) \geq w(p_{nT_0}, q_{nT_0})$ and he plays an optimal strategy in the non-revealing game of length T_0 at (p_{nT_0}, q_{nT_0}), or (2) $\hat{v}(p_{nT_0}, q_{nT_0}) < w(p_{nT_0}, q_{nT_0})$, and he can send a random message to Player 2 in order to drive his beliefs at some points $\tilde{p}_{nT_0}$ such that $\hat{v}(\tilde{p}_{nT_0}, q_{nT_0}) \geq w(\tilde{p}_{nT_0}, q_{nT_0})$ and there he plays as in case (1). This strategy would defend $w(p_{nT_0}, q_{nT_0})$ up to an error of ε on the n-th block of size T_0 if Player 2 was not using his information on the recurrence classes of the chain $(l_t)_t$ on this block. In order to bound the error due to this approximation, we have to replace the error term based on the L_1-variation of the process $(q_t)_{t\geq 1}$ (which is no more a martingale here) by the L_1-variation of an auxiliary martingale representing the beliefs of Player 1 over the recurrence classes of the Markov chain $(l_t)_{t\geq 1}$ (note however that we obtain a less precise bound since the method developed for games with incomplete information does not apply here). Then, we prove that the expected value of $w(p_{nT_0}, q_{nT_0})$ is always greater or equal to $w(p,q)$ using similar tools as in [14] and also that w is balanced. We obtain $\text{liminf}_n v_n \geq \sup\{w | w \in \mathcal{W}^-(\hat{v}), \text{balanced}\}$, and by symmetry $\text{limsup}_n v_n \leq \inf\{w | w \in \mathcal{W}^+(\hat{v}), \text{balanced}\}$.

Then it remains to show the equality

$$\sup\{w | w \in \mathcal{W}^-(\hat{v}), \text{balanced}\} = \inf\{w | w \in \mathcal{W}^+(\hat{v}), \text{balanced}\},$$

by reducing the problem to the equality showed by Mertens and Zamir. The conclusion follows.

5 Non revealing games

Let σ in Σ be a strategy of Player 1, i.e. $\sigma = (\sigma_T)_{T\geq 1}$, with $\sigma_T : (K \times I \times J)^{T-1} \times K \longrightarrow \Delta(I)$ for all T. For each finite history h_T in $(I \times J)^T$, we say that h_T is compatible with (p, σ) if for some τ and q we have $I\!P_{p,q,\sigma,\tau}(h_T) > 0$. In this case we define $p_T(p,\sigma)(h_T)$ in $\Delta(K)$ as the law of the state k_{T+1} knowing that: the initial state k_1 is selected according to p, Player 1 uses σ and h_T has been played at the first T stages. It does not depend on the last move of Player 2 in h_T, and for all k in K, q in $\Delta(L)$ and strategy τ of Player 2 such that $I\!P_{p,q,\sigma,\tau}(h_T) > 0$, we have:

$$p^k_T(p,\sigma)(h_T) = I\!P_{p,q,\sigma,\tau}(k_{T+1} = k | h_T).$$

If h_T is not compatible with (p,σ), we define $p_T(p,\sigma)(h_T)$ arbitrarily in $\Delta(K)$. For $T = 0$, $p_0(p,\sigma) = p$ is the law of the first state k_1.

Definition 5.1. *The relevant information of Player 2 after h_T has been played is:*

$$\hat{p}_T(p,\sigma)(h_T) = p_T(p,\sigma)(h_T)B.$$

For each T, we denote by $\mathcal{H}_T$ the σ-algebra on the set of plays Ω generated by the projection on $(I \times J)^T$ giving the first T moves.

Lemma 5.2. *For any strategy pair (σ,τ) in $\Sigma \times \mathcal{T}$, the process $(\hat{p}_T(p,\sigma))_{T\geq 0}$ is a $(\mathcal{H}_T)_{T\geq 0}$-martingale with respect to $I\!P_{p,q,\sigma,\tau}$.*

Proof. Assume (σ,τ) is played, the initial probabilities on $\Delta(K)$ and $\Delta(L)$ being p and q. Fix $T \geq 0$ and $h_T = (i_1, j_1, ..., i_T, j_T) \in (I \times J)^T$. Given (i_{T+1}, j_{T+1}) in $I \times J$, we write $h_{T+1} = (i_1, ..., j_T, i_{T+1}, j_{T+1})$ and denote by $r_T(p,\sigma)(h_{T+1})$ the conditional probability on the state k_{T+1} given that h_{T+1} has been played at the first $T+1$ stages. We have $r_T(p,\sigma)(h_{T+1})M = p_{T+1}(p,\sigma)(h_{T+1})$, and $I\!E_{p,q,\sigma,\tau}(r_T(p,\sigma)|h_T) = p_T(p,\sigma)(h_T)$. Now,

$$\begin{aligned}
I\!E_{p,q,\sigma,\tau}(\hat{p}_{T+1}(p,\sigma)|h_T) &= I\!E_{p,q,\sigma,\tau}(p_{T+1}(p,\sigma)B|h_T) \\
&= I\!E_{p,q,\sigma,\tau}(r_T(p,\sigma)MB|h_T) \\
&= I\!E_{p,q,\sigma,\tau}(r_T(p,\sigma)B|h_T) \\
&= p_T(p,\sigma)(h_T)B \\
&= \hat{p}_T(p,\sigma)(h_T).
\end{aligned}$$

□

Definition 5.3. *A strategy σ in Σ is called non revealing at p if for all τ and q, the martingale $(\hat{p}_T(p,\sigma))_{T\geq 0}$ is $I\!P_{p,q,\sigma,\tau}$-almost surely constant. The set of NR strategies of Player 1 at p is denoted $\hat{\Sigma}(p)$.*

We will give an alternative definition of non revealing strategies.

Definition 5.4.

$$\begin{aligned} NR(p) &= \{x \in \Delta(I)^K, \forall i \in I \text{ } s.t. \text{ } x(p)(i) > 0, \;\; \vec{p}(x,i)B = pB\} \\ &= \{x \in \Delta(I)^K, \forall i \in I, (p^k x^k(i))_{k\in K} B = x(p)(i)\, pB\}. \end{aligned}$$

Given σ in Σ and an history $h_T \in (I\times J)^T$ which is compatible with (p,σ), for each i in I and k in K, we denote by $x^k(p,\sigma)(h_T)(i)$ the probability $I\!P_{p,q,\sigma,\tau}(i_{T+1} = i | k_{T+1} = k, h_T)$ that Player 1 plays i after h_T if the next state is k. This probability does not depend on (q,τ). The vector $x(p,\sigma)(h_T) = (x^k(p,\sigma)(h_T)(i))_{k,i}$ is viewed as en element of $\Delta(I)^K$.

Lemma 5.5. *A strategy σ in Σ is non revealing at p if and only if for all $T\geq 0$ and history $h_T \in (I\times J)^T$ compatible with (p,σ), the vector $x(p,\sigma)(h_T)$ belongs to $NR(p_T(p,\sigma)(h_T))$.*

Proof. The proof is similar to the proof of Proposition 6.5 in [20]. Fix p and σ. Consider $T\geq 0$ and a history h_T in $(I\times J)^T$ compatible with (p,σ), and let (q,τ) be such that $I\!P_{p,q,\sigma,\tau}(h_T) > 0$. For simplicity we write $x = x(p,\sigma)(h_T)$ in $\Delta(I)^K$, and $p_T = p_T(p,\sigma)(h_T) \in \Delta(K)$.

$$\begin{aligned} \forall i \in I, \; I\!P_{p,q,\sigma,\tau}(i_{T+1} = i|h_T) &= \sum_{k\in K} I\!P_{p,q,\sigma,\tau}(k_{T+1} = k|h_T) x^k(i), \\ &= \sum_{k\in K} p_T^k x^k(i), \\ &= x(p_T)(i). \end{aligned}$$

Consider now a history $h_{T+1} = (h_T, i_{T+1}, j_{T+1})$ extending h_T such that $I\!P_{p,q,\sigma,\tau}(h_{T+1}) > 0$. We have $(I\!P_{p,q,\sigma,\tau}(k_{T+1} = k|h_{T+1}))_{k\in K} = \vec{p}_T(x, i_{T+1})$, so $p_{T+1}(p,\sigma)(h_{T+1}) = \vec{p}(x,i_{T+1})M$. Consequently:

$$\hat{p}_{T+1}(p,\sigma)(h_{T+1}) = \vec{p}_T(x, i_{T+1})B. \tag{2}$$

We now conclude the proof of the lemma.

$\Longrightarrow$ Assume that σ is non revealing at p, and consider T and h_T in $(I\times J)^T$ such that h_T is compatible with (p,σ). Write $p_T = p_T(p,\sigma)(h_T)$ and $x = x(p,\sigma)(h_T)$. If $x(p_T)(i_{T+1}) > 0$, then (h_T, i_{T+1}) is compatible with (p,σ) and by assumption $\hat{p}_{T+1}(p,\sigma)(h_T, i_{T+1}) = \hat{p}_T(p,\sigma)(h_T)$. We have $\hat{p}_{T+1}(p,\sigma)(h_T,i_{T+1}) = \vec{p}_T(x,i_{T+1})B$ by equation (2), and by definition $\hat{p}_T(p,\sigma)(h_T) = p_T B$. We obtain $\vec{p}_T(x,i_{T+1})B = p_T B$, so $x \in NR(p_T)$.

$\Longleftarrow$ Assume that for all $T\geq 0$ and history $h_T \in (I\times J)^T$ compatible with (p,σ), the vector $x(p,\sigma)(h_T)$ belongs to $NR(p_T(p,\sigma)(h_T))$. Consider $T\geq 0$, and a history $h_{T+1} = (i_1, j_1, ..., i_{T+1}, j_{T+1})$ compatible with (p,σ). Write $h_T = (i_1, ... j_T)$, $p_T = p_T(p,\sigma)(h_T)$, and $x = x(p,\sigma)(h_T)$. By equation (2), we have $\hat{p}_{T+1}(p,\sigma)(h_{T+1}) = \vec{p}_T(x,i_{T+1})B$. We have $x(p_T)(i_{T+1}) > 0$ because h_{T+1} is compatible with (p,σ), so by assumption, $\vec{p}_T(x,i_{T+1})B = p_T B$, and by definition $\hat{p}(p,\sigma)(h_T) = p_T B$. We obtain $\hat{p}_{T+1}(p,\sigma)(h_{T+1}) = \hat{p}_T(p,\sigma)(h_T)$. □

We also have a splitting lemma with non revealing strategies.

Lemma 5.6. *Consider a convex combination $p = \sum_{s\in S} \alpha_s p_s$ in $\Delta(K)$ having the property that $p_s B = pB$ for all s. Consider for each S a non revealing strategy σ_s in $\hat{\Sigma}(p_s)$. There exists σ in $\hat{\Sigma}(p)$ such that:*

$$\forall q \in \Delta(L), \forall \tau \in \mathcal{T}, \;\; I\!P_{p,q,\sigma,\tau} = \sum_{s\in S} \alpha_s I\!P_{p_s,q,\sigma_s,\tau}.$$

The proof is similar to the proof of Lemma 6.6 in [20], with σ being defined via the splitting procedure: observe the first state k_1 in K, then choose s according to the probability $\alpha_s p_s^{k_1}/p^{k_1}$, and play according to σ_s. Notice that the equality $I\!P_{p,q,\sigma,\tau} = \sum_{s\in S} \alpha_s I\!P_{p_s,q,\sigma_s,\tau}$ is valid for any, revealing or non revealing, strategy τ of Player 2. See also Lemma 6.6 where we study more precise properties of this splitting procedure.

Non revealing strategies of Player 2 at q are defined similarly, the set of such strategies is denoted $\hat{\mathcal{T}}(q)$.

Definition 5.7. *The T-stage non revealing game at (p,q) is the T-stage game where the strategy sets are restricted to $\hat{\Sigma}(p)$ and $\hat{\mathcal{T}}(q)$. It is denoted by $\hat{\Gamma}_T(p,q)$.*

Define $\hat{\Sigma}_T(p)$ as the set of T-stage non revealing strategies of Player 1 at p. Formally, $\hat{\Sigma}_T(p)$ is the projection of $\hat{\Sigma}(p)$ over the (compact) set of T-stage strategies of Player 1. Similarly, $\hat{\mathcal{T}}_T(q)$ denotes the set of T-stage non revealing strategies of Player 2 at q. The game $\hat{\Gamma}_T(p,q)$ can equivalently be seen as the zero-sum game $(\hat{\Sigma}_T(p), \hat{\mathcal{T}}_T(q), \gamma_T^{p,q})$.

Proposition 5.8. *For all $T \geq 1$, the T-stage non revealing game at (p,q) has a value denoted by $\hat{v}_T(p,q)$. For each q in $\Delta(L)$ and p^* in P^*, $(p \mapsto \hat{v}_T(p,q))$ is concave on $B^{-1}(p^*)$, and similarly for each p in $\Delta(K)$ and q^* in Q^*, $(q \mapsto \hat{v}_T(p,q))$ is convex on $C^{-1}(q^*)$.*

Moreover, for all $T \geq 0$ we have the recursive formula:

$$\begin{aligned}(T+1)\hat{v}_{T+1}(p,q) &= \max_{x\in NR(p)} \min_{y\in NR(q)} \left(G(p,q,x,y) + T \sum_{i\in I, j\in J} x(p)(i) y(q)(j) \hat{v}_T(\vec{p}(x,i)M, \vec{q}(y,j)N) \right) \\ &= \min_{y\in NR(q)} \max_{x\in NR(p)} \left(G(p,q,x,y) + T \sum_{i\in I, j\in J} x(p)(i) y(q)(j) \hat{v}_T(\vec{p}(x,i)M, \vec{q}(y,j)N) \right).\end{aligned}$$

Proof: Consider a sequence $(p_a, \sigma_a)_{a\geq 1}$ converging to (p,σ) where $\sigma_a \in \hat{\Sigma}_T(p_a)$ for each a. For each history h_T in $(I \times J)^T$ and τ in $\mathcal{T}$ such that $I\!P_{p,q,\sigma,\tau}(h_T) > 0$, the conditional $(I\!P_{p_a,q,\sigma_a,\tau}(k_{T+1} = k|h_T))_{k\in K}$ converges to $(I\!P_{p,q,\sigma,\tau}(k_{T+1} = k|h_T))_{k\in K}$, so we can pass to the limit in the equality $p_T(p_a,\sigma_a)(h_T)B = p_a B$ to obtain that σ is a T-stage non revealing strategy of Player 1 at p. As a consequence, $\{(p,\sigma), \sigma \in \hat{\Sigma}_T(p)\}$ is a closed subset of $\Delta(K) \times \Sigma_T(p)$, hence is compact.

By Lemma 5.6, we obtain that the game $(\hat{\Sigma}_T(p), \hat{\mathcal{T}}_T(q), \gamma_T^{p,q})$ is affine-like: for all $\sigma', \sigma'' \in \hat{\Sigma}_T(p)$, for all $\alpha \in [0,1]$, there exists σ in $\hat{\Sigma}_T(p)$ such that for all τ, $\gamma_T^{p,q}(\sigma,\tau) = \alpha\gamma_T^{p,q}(\sigma',\tau) + (1-\alpha)\gamma_T^{p,q}(\sigma'',\tau)$.

Summing up, we can use a minmax theorem due to Fan (1953 see Proposition A.13 in [25]) to obtain that that the T-stage non revealing game at (p,q) has a value $\hat{v}_T(p,q)$, and that both players have optimal strategies.

Since $\{(p,\sigma), \sigma \in \hat{\Sigma}_T(p)\}$ is compact, $\hat{v}_T(p,q)$ is upper semi-continuous in p, and by symmetry it is lower semi-continuous in q. Lemma 5.6 gives that $\hat{v}_T(p,q)$ is concave in p on each set $B^{-1}(p^*)$, and convex in q on each set $C^{-1}(q^*)$.

We now prove the recursive formula in a standard way. Fix $T \geq 0$ and (p,q) in $\Delta(K) \times \Delta(L)$. We consider the zero-sum game $A_{T+1}(p,q) = (NR(p), NR(q), f)$, where $f(x,y) = \frac{1}{T+1}G(p,q,x,y) + \frac{T}{T+1}\sum_{i\in I, j\in J} x(p)(i)y(q)(j)\hat{v}_T(\vec{p}(x,i)M, \vec{q}(y,j)N)$. The strategy sets $NR(p)$ and $NR(q)$ are convex compact, and one can prove (as in the proof of Proposition 7.4 in [20]) that f is concave s.c.s. in x and convex l.s.c. in y. By Sion's minmax Theorem, $A_{T+1}(p,q)$ has a value that we denote by $a_{T+1}(p,q)$.

Consider now an optimal strategy τ of Player 2 in the non revealing game $\hat{\Gamma}_{T+1}(p,q)$. Since τ is non revealing at q, the first stage strategy τ_1 lies in $NR(q)$. Define x in $NR(p)$ which is a best response against $y = \tau_1$ in $A_{T+1}(p,q)$, and a strategy σ of Player 1 which: plays according to x at stage 1, and from stage 2 on play an optimal strategy in the game $\hat{\Gamma}_T(\vec{p}(x,i)M, \vec{q}(y,j)N)$, where i and j are the moves played at stage 1 (with $\vec{q}(y,j)$ defined arbitrarily if $y(q)(j) = 0$). Using Lemma 5.5, one can show that the strategy σ is non revealing at p, and the payoff $\gamma_T^{p,q}(\sigma,\tau)$ is at least $f(x,y)$, so at least $a_{T+1}(p,q)$. Since τ was optimal, we deduce that $a_{T+1}(p,q) \leq \hat{v}_{T+1}(p,q)$. Exchanging the roles of the players, we get $a_{T+1}(p,q) = \hat{v}_{T+1}(p,q)$ and the recursive formula is proved. This ends the proof of Proposition 5.8. □

We now prove a proposition allowing to consider smaller sets of strategies.

Definition 5.9. *Let $\overline{\mathcal{T}}$ denote the subset of strategies of Player 2 which for all $T \geq 0$, depend at stage $T+1$ only on the past history $h_T \in (I \times J)^T$ and on the current state l_{T+1}. Define $\overline{\Sigma}$ similarly.*

Proposition 5.10. *For all $(p,q) \in \Delta(K) \times \Delta(L)$, for all $\tau \in \mathcal{T}$, there exists a strategy $\overline{\tau} \in \overline{\mathcal{T}}$ such that for all $T \geq 1$ and $\sigma \in \Sigma$, $\gamma_T^{p,q}(\sigma,\tau) = \gamma_T^{p,q}(\sigma,\overline{\tau})$. Moreover, if τ is non-revealing at q, then $\overline{\tau}$ is also non-revealing at q.*

As a corollary, in $\Gamma_T(p,q)$ and in the non revealing game $\hat{\Gamma}_T(p,q)$ the players have optimal strategies in $\overline{\Sigma}$ and $\overline{\mathcal{T}}$.

Although it is intuitive that Player 2 does not need to remember the whole sequence of states $(l_1, ..., l_t, ...)$, the proof of the proposition requires to be precise.

Proof of Proposition 5.10. The strategy $\overline{\tau}$ proceeds as follows: for $t \geq 0$, at stage $t+1$, Player 2 does not remind the past states $(l_1, .., l_t)$, but using the history h_t, he can generate a virtual sequence $(\overline{l}_1, .., \overline{l}_t)$ using the conditional law of $(l_1, .., l_t)$ given (h_t, l_{t+1}) under $\mathbb{P}_{p,q,\sigma,\tau}$ (which does not depend on σ). He selects then at stage $t+1$ an action j_{t+1} with the probability $\tau(h_t, \overline{l}_1, .., \overline{l}_t, l_{t+1})$. Formally, we have

$$\overline{\tau}(h_t)(l_{t+1}) = \mathbb{E}_{p,q,\sigma,\tau}[\tau(h_t, l_1, ..., l_t, l_{t+1}) | h_t, l_{t+1}].$$

We will prove that for all $t \geq 1$, (h_t, l_t, k_t) has the same distribution under $I\!\!P_{p,q,\sigma,\tau}$ and under $I\!\!P_{p,q,\sigma,\overline{\tau}}$, i.e. for all $h \in H_t$, for all $(k,l) \in K \times L$,

$$I\!\!P_{p,q,\sigma,\tau}(h_t = h, l_t = l, k_t = k) = I\!\!P_{p,q,\sigma,\overline{\tau}}(h_t = h, l_t = l, k_t = k). \tag{3}$$

Let us proceed by induction on t. For $t = 1$, the property is obviously true since $\tau_1 = \overline{\tau}_1$. Assume that the property is true for $t \geq 1$. At first, the conditional distribution of l_{t+1} given (h_t, k_t, l_t) is $M(l_t, .)$ under $I\!\!P_{p,q,\sigma,\tau}$ and under $I\!\!P_{p,q,\sigma,\overline{\tau}}$. We deduce that the law of (h_t, l_{t+1}) is the same under both probabilities. By construction the conditional law of j_{t+1} given (h_t, l_{t+1}) is the same under $I\!\!P_{p,q,\sigma,\tau}$ and under $I\!\!P_{p,q,\sigma,\overline{\tau}}$, which implies that the conditional law of (l_{t+1}, j_{t+1}) given h_t is also the same. Using that (k_{t+1}, i_{t+1}) and (l_{t+1}, j_{t+1}) are conditionally independent given h_t (under both distributions) and that the conditional law of (k_{t+1}, i_{t+1}) given h_t does not depend on τ, the conditional distribution of $(k_{t+1}, l_{t+1,}, i_{t+1}, j_{t+1})$ given h_t is the same under $I\!\!P_{p,q,\sigma,\tau}$ and under $I\!\!P_{p,q,\sigma,\overline{\tau}}$. We deduce that the law of $(h_{t+1}, k_{t+1}, l_{t+1})$ is the same under both probability distributions which concludes the proof of (3). Going back to the main proof, we obtain

$$\begin{aligned}
\gamma_T^{p,q}(\sigma, \tau) &= \sum_{t=0}^{T-1} I\!\!E_{p,q,\sigma,\tau}[I\!\!E_{p,q,\sigma,\tau}[g(k_{t+1}, l_{t+1}, i_{t+1}, j_{t+1})|h_t]] \\
&= \sum_{t=0}^{T-1} I\!\!E_{p,q,\sigma,\overline{\tau}}[I\!\!E_{p,q,\sigma,\overline{\tau}}[g(k_{t+1}, l_{t+1}, i_{t+1}, j_{t+1})|h_t]] \\
&= \gamma_T^{p,q}(\sigma, \overline{\tau})
\end{aligned}$$

Moreover, since we proved that the law of (h_t, l_{t+1}) is the same under $I\!\!P_{p,q,\sigma,\tau}$ and under $I\!\!P_{p,q,\sigma,\overline{\tau}}$, this implies that $q_t(q, \tau)$ and $q_t(q, \overline{\tau})$ also have the same law, which proves the last assertion.

Remark 5.11. Note that in the above construction, the laws $I\!\!P_{p,q,\sigma,\tau}$ and $I\!\!P_{p,q,\sigma,\overline{\tau}}$ do not coincide on the whole set Ω but generate the same payoffs. The reader may convince himself by considering the following example:

$$L = \{1, 2\}, \ N = \begin{pmatrix} 1/3 & 2/3 \\ 2/3 & 1/3 \end{pmatrix}, \ q = (1/2, 1/2), \ J = \{1, 2\},$$

and τ is defined by: play 1 at stage 1 and then at stage $t \geq 2$, play 1 if $l_t = l_{t-1}$ and 2 otherwise. A direct computation leads to

$$I\!\!P_{p,q,\sigma,\tau}(l_1 = l_2 = 1, j_2 = 1) = 1/6, \quad I\!\!P_{p,q,\sigma,\overline{\tau}}(l_1 = l_2 = 1, j_2 = 1) = 1/18.$$

Remark 5.12. In the above proposition, we made use of some standard properties of $I\!\!P_{p,q,\sigma,\tau}$, namely for all $t \geq 1$: the conditional law of $(l_1, .., l_t)$ given (h_t, l_{t+1}) does not depend on σ, the conditional law of (k_{t+1}, i_{t+1}) given h_t does not depend on τ, (k_{t+1}, i_{t+1}) and (l_{t+1}, j_{t+1}) are conditionally independent given h_t. All these properties can be obtained by direct algebraic manipulations, the details are left to the reader.

The next proposition is a major point of our proof.

Proposition 5.13. *For all $T \geq 1$, $\hat{v}_T$ is 3-Lipschitz on $\Delta(K) \times \Delta(L)$.*

The proof of Proposition 5.13 being quite technical, it is postponed to section 7. The crucial consequence is to obtain uniform equi-continuity of the family of functions $(\hat{v}_T)_{T\geq 1}$. In Renault [20], the functions $(\hat{v}_T)_{T\geq 1}$ where only proved to be u.s.c. and this property would not have been sufficient here, since the proof of our main Theorem 6.1 relies on uniform convergence of the $(\hat{v}_T)_{T\geq 1}$ to their limit. The main point is to introduce a function $S(p,p')$ (see Definition 7.2) which will play the role of a metric and with respect to which all the functions $(\hat{v}_T)_{T\geq 1}$ are "1-Lipschitz". However, we do not know if the functions $(\hat{v}_T)_{T\geq 1}$ are 1-Lipschitz with respect to $\|.\|$ or if there exists a real metric having this property.

Before proving that the non revealing values $\hat{v}_T$ converge when T goes to infinity, we now present a definition and a lemma, allowing to concatenate non revealing strategies defined on blocks.

Definition 5.14. *Given $T_0, T_1 \geq 1$ and a strategy $\sigma \in \Sigma$ such that for all $1 \leq t \leq T_1$, σ_{T_0+t} does not depend on $(k_1, ..., k_{T_0})$, we define for all $h_{T_0} \in H_{T_0}$ a T_1-stage strategy $\sigma(h_{T_0})$. This strategy $\sigma(h_{T_0})$ plays, after a history $(h'_{t-1}, k'_1, ..., k'_t)$, what σ plays at stage T_0+t after $((h_0, h'_{t-1}), k'_1, ..., k'_t)$. Formally,*

$$\sigma(h_{T_0})_t(h'_{t-1}, k'_1, ..., k'_t) = \sigma_{T_0+t}((h_{T_0}, h'_{t-1}), k'_1, ..., k'_t), \forall 1 \leq t \leq T_1, \forall h'_t \in H_{t-1}, \forall (k'_1, .., k'_t) \in K^t.$$

Lemma 5.15. *Consider $T_0, T_1 \in \mathbb{N}^*$ and strategies $(\sigma, \tau) \in \Sigma \times \mathcal{T}$ such that for all $1 \leq t \leq T_1$, σ_{T_0+t} and τ_{T_0+t} do not depend on, respectively, $(k_1, ..., k_{T_0})$ and $(l_1, .., l_{T_0})$. The conditional law of $(k_{T_0+t}, l_{T_0+t}, i_{T_0+t}, j_{T_0+t})_{t=1,..,T_1}$ given h_{T_0} under $\mathbb{P}_{p,q,\sigma,\tau}$ is precisely $\mathbb{P}_{p_{T_0}, q_{T_0}, \sigma(h_{T_0}), \tau(h_{T_0})}$.*

The proof follows easily from a direct computation.

Remark 5.16. *Note that if $\tau \in \overline{\mathcal{T}}$, then the above condition on τ holds for all T_0, T_1. We will use the following consequences in the sequel, at first*

$$\mathbb{E}_{p,q,\sigma,\tau}[\sum_{t=1}^{T_1} g(k_{T_0+t}, l_{T_0+t}, i_{T_0+t}, j_{T_0+t})] = \mathbb{E}_{p,q,\sigma,\tau}[\gamma_{T_1}^{p_{T_0}, q_{T_0}}(\sigma(h_{T_0}), \tau(h_{T_0}))],$$

and for all $t = 1, ..., T_1$ and any continuous function f defined on $\Delta(K) \times \Delta(L)$, for all h_{T_0} in $(I \times J)^{T_0}$,

$$\mathbb{E}_{p,q,\sigma,\tau}[f(p_{T_0+t}, q_{T_0+t})|h_{T_0}] = \mathbb{E}_{p_{T_0}, q_{T_0}, \sigma(h_{T_0}), \tau(h_{T_0})}[f(p_t, q_t)],$$

where $p_{T_0} = p_{T_0}(p, \sigma)(h_{T_0})$ and $q_{T_0} = q_{T_0}(q, \tau)(h_{T_0})$.

Proposition 5.17. *$\hat{v}(p,q) = \lim_{T\to\infty} \hat{v}_T(p,q)$ exists and is balanced.*

Proof. Let $(p,q) \in \Delta(K) \times \Delta(L)$, $p^* = pB$, $q^* = qC$ and $\varepsilon > 0$. Choose $T_0, T_1 \in \mathbb{N}^*$ such that

$$\|M^{T_0} - B\| \leq \varepsilon, \quad \|N^{T_0} - C\| \leq \varepsilon, \quad \frac{T_0}{T_1} \leq \varepsilon,$$

$$\text{and} \quad \hat{v}_{T_1}(p^*, q^*) \geq \text{limsup}_T \hat{v}_T(p^*, q^*) - \varepsilon.$$

Given $N \in \mathbb{N}^*$ and an optimal strategy $\tau \in \overline{\mathcal{T}}$ of Player 2 in the game $\hat{\Gamma}_{N(T_0+T_1)}(p,q)$, let us construct the strategy σ as follows. For $n = 0$ to $N-1$, during stages $t = n(T_0+T_1)+1$ to $t = n(T_0+T_1)+T_0$, play a fixed action $i_0 \in I$. During the next T_1 stages, play an optimal strategy in the game $\hat{\Gamma}_{T_1}(p_{n(T_0+T_1)+T_0}(p,\sigma), q_{n(T_0+T_1)+T_0}(q,\tau))$. The payoff can be written as

$$\begin{aligned}
N(T_0+T_1)\gamma^{p,q}_{N(T_0+T_1)}(\sigma,\tau) &= \sum_{n=0}^{N-1}\left(\sum_{m=n(T_0+T_1)+1}^{n(T_0+T_1)+T_0} \mathbb{E}_{p,q,\sigma,\tau}[g(k_m,l_m,i_m,j_m)]\right.\\
&\qquad \left.+ \sum_{m=n(T_0+T_1)+T_0+1}^{(n+1)(T_0+T_1)} \mathbb{E}_{p,q,\sigma,\tau}[g(k_m,l_m,i_m,j_m)]\right)\\
&\geq \sum_{n=0}^{N-1}\left(-T_0 + T_1\mathbb{E}_{p,q,\sigma,\tau}[\gamma_{T_1}^{p_{n(T_0+T_1)}(p,\sigma),q_{n(T_0+T_1)}(q,\tau)}(\sigma(h_{n(T_0+T_1)}),\tau(h_{n(T_0+T_1)}))]\right)\\
&\geq \sum_{n=0}^{N-1}\left(-T_0 + T_1\mathbb{E}_{p,q,\sigma,\tau}[\hat{v}_{T_1}(p_{n(T_0+T_1)+T_0}(p,\sigma), q_{n(T_0+T_1)+T_0}(q,\tau))]\right)\\
&\geq \sum_{n=0}^{N-1}\left(-T_0 + T_1\mathbb{E}_{p,q,\sigma,\tau}[\hat{v}_{T_1}(p_{n(T_0+T_1)}(p,\sigma)M^{T_0}, q_{n(T_0+T_1)}(q,\tau)N^{T_0})]\right)\\
&\geq \sum_{n=0}^{N-1}\left(-T_0 + T_1(\hat{v}_{T_1}(p^*,q^*)-6\varepsilon)\right).
\end{aligned}$$

The first inequality follows from Lemma 5.15 and the fact that g is valued in $[-1,1]$, the second follows from the definition of σ. Note then that

$$p_{n(T_0+T_1)+T_0}(p,\sigma) = p_{n(T_0+T_1)}(p,\sigma)M^{T_0}$$

since the strategy σ is independent of the state variable during the block of size T_0 and that

$$\mathbb{E}_{p,q,\sigma,\tau}[q_{n(T_0+T_1)+T_0}(q,\tau)|h_{n(T_0+T_1)}] = q_{n(T_0+T_1)}(q,\tau)N^{T_0}.$$

The third inequality is obtained by taking conditional expectation with respect to $h_{n(T_0+T_1)}$ and using that $\hat{v}_{T_1}$ is convex with respect to the second variable on $C^{-1}(q^*)$. The last inequality follows directly from the properties of T_0 and the fact that $\hat{v}_{T_1}$ is 3-Lipschitz. We deduce that for all $N \in \mathbb{N}^*$

$$\hat{v}_{N(T_0+T_1)}(p,q) \geq \frac{-T_0 + T_1(\hat{v}_{T_1}(p^*,q^*)-6\varepsilon)}{(T_0+T_1)} \geq \hat{v}_{T_1}(p^*,q^*) - 8\varepsilon \geq \mathrm{limsup}_T \hat{v}_T(p^*,q^*) - 9\varepsilon.$$

It follows that

$$\mathrm{liminf}_N \hat{v}_{N(T_0+T_1)}(p,q) \geq \mathrm{limsup}_T \hat{v}_T(p^*,q^*) - 9\varepsilon.$$

Using that $\|\hat{v}_T - \hat{v}_{T+T'}\|_\infty \leq \frac{2T'}{T}$, this implies

$$\mathrm{liminf}_T \hat{v}_T(p,q) \geq \mathrm{limsup}_T \hat{v}_T(p^*,q^*) - 9\varepsilon$$

and finally $\mathrm{liminf}_T \hat{v}_T(p,q) \geq \mathrm{limsup}_T \hat{v}_T(p^*,q^*)$. Inverting the role of K and L leads to $\mathrm{limsup}_T \hat{v}_T(p,q) \leq \mathrm{liminf}_T \hat{v}_T(p^*,q^*)$, and we conclude that $\hat{v}(p,q) = \lim_{T\to\infty} \hat{v}_T(p,q) = \lim_T \hat{v}_T(p^*,q^*)$ exists and is a balanced function. $\square$

6 Back to the original game and the Mertens-Zamir system

Recall that $\mathcal{W}$ denotes the set of continuous functions $w : \Delta(K) \times \Delta(L) \longrightarrow [-1, 1]$.

Theorem 6.1. *$v = \lim_T v_T$ exists and can be characterized by:*

$$v = \inf \left\{ \begin{matrix} w \in \mathcal{W}, w \text{ is } I\text{-concave, balanced and} \\ \forall (p,q) \in \Delta(K) \times \Delta(L),\ w(p,q) \geq \mathrm{vex}_{II}\mathrm{Max}(w, \hat{v})(p,q) \end{matrix} \right\},$$
$$= \sup \left\{ \begin{matrix} w \in \mathcal{W}, w \text{ is } II\text{-convex, balanced and} \\ \forall (p,q) \in \Delta(K) \times \Delta(L),\ w(p,q) \leq \mathrm{cav}_{I}\mathrm{Min}(w, \hat{v})(p,q) \end{matrix} \right\}.$$

Before we prove Theorem 6.1, we will prove a proposition and two lemmas. We first show that the comparison result of Mertens and Zamir can be extended to our case.

Proposition 6.2. *If we define*

$$\overline{v} = \inf \left\{ \begin{matrix} w \in \mathcal{W}, w \text{ is } I\text{-concave, balanced and} \\ \forall (p,q) \in \Delta(K) \times \Delta(L),\ w(p,q) \geq \mathrm{vex}_{II}\mathrm{Max}(w, \hat{v})(p,q) \end{matrix} \right\},$$

$$\underline{v} = \sup \left\{ \begin{matrix} w \in \mathcal{W}, w \text{ is } II\text{-convex, balanced and} \\ \forall (p,q) \in \Delta(K) \times \Delta(L),\ w(p,q) \leq \mathrm{cav}_{I}\mathrm{Min}(w, \hat{v})(p,q) \end{matrix} \right\},$$

then $\overline{v} = \underline{v}$.

Proof. Consider any balanced function w in $\mathcal{W}$. One may identify w with its restriction w^* to the product of simplices $P^* \times Q^*$. I-concavity and II-convexity on $\Delta(K) \times \Delta(L)$ of w is equivalent to I-concavity and II-convexity of w^* on $P^* \times Q^*$ (using that B and C are linear). We will prove that $\mathrm{cav}_I(w)$ is balanced and that $(\mathrm{cav}_I w)^* = \mathrm{cav}_I(w^*)$.

Fix (p,q) be in $\Delta(K) \times \Delta(L)$, and define $p^* = pB$ and $q^* = qB$. Recall that

$$\mathrm{cav}_I w(p,q) = \sup\{\sum_{m=1}^{N} \alpha_m w(p_m, q) \mid N \geq 1, \forall m = 1,..,N,\ \alpha_m \geq 0,\ \sum_{m=1}^{N} \alpha_m = 1,\ \sum_{m=1}^{N} \alpha_m p_m = p\}.$$

Given $(\alpha_m, p_m)_{m=1,..,N}$ as above, for all m, $w(p_m, q) = w(p_m B, q^*)$, and $\sum_{m=1}^{N} \alpha_m p_m B = p^*$, which implies

$$\sum_{m=1}^{N} \alpha_m w(p_m, q) = \sum_{m=1}^{N} \alpha_m w(p_m B, q^*) \leq \mathrm{cav}_I(w^*)(p^*, q^*),$$

and therefore $\mathrm{cav}_I w(p,q) \leq \mathrm{cav}_I(w^*)(p^*, q^*)$.

Define now the affine map f from P^* to $\Delta(K)$, depending on p, by:

$$\forall p' \in P^*, \forall r = 1,..,r_M, \forall k \in K(r),\ f(p')^k = \begin{cases} (\sum_{s \in K(r)} p'^s) & \frac{p^k}{\sum_{s \in K(r)} p^s} \quad \text{if} \quad \sum_{s \in K(r)} p^s > 0 \\ (\sum_{s \in K(r)} p'^s) & p^*(r)^k \quad \text{if} \quad \sum_{s \in K(r)} p^s = 0 \end{cases}.$$

We have $f(p^*) = p$. Moreover, for any recurrence class $K(r)$, we have $\sum_{k\in K(r)} f(p')^k = \sum_{s\in K(r)} p'^s$. This implies that for all $p' \in P^*$, $f(p')B = p'$. Consider now $(\alpha_m, p^*_m)_{m=1,..,N}$ such that

$$\forall m = 1, ..., N,\ \alpha_m \geq 0, p^*_m \in P^*,\ \sum_{m=1}^{N} \alpha_m = 1,\ \sum_{m=1}^{N} \alpha_m p^*_m = p^*.$$

We have by construction $\sum_{m=1}^{N} \alpha_m f(p^*_m) = p$ and $w(p^*_m, q^*) = w(f(p^*_m), q)$ which implies

$$\sum_{m=1}^{N} \alpha_m w(p^*_m, q^*) = \sum_{m=1}^{N} \alpha_m w(f(p^*_m), q) \leq \mathrm{cav}_I w(p, q),$$

and therefore $\mathrm{cav}_I(w^*)(p^*, q^*) \leq \mathrm{cav}_I w(p, q)$. We conclude that:

$$\forall (p, q) \in \Delta(K) \times \Delta(L),\ \mathrm{cav}_I w(p, q) = \mathrm{cav}_I(w^*)(p^*, q^*).$$

Consequently, $\mathrm{cav}_I(w)$ is balanced and $(\mathrm{cav}_I w)^* = \mathrm{cav}_I(w^*)$.
Since $\hat{v}$ is also balanced, we deduce that :

$$\underline{v}^* = \sup \left\{ \begin{matrix} w : P^* \times Q^* \longrightarrow [-1, 1], w \text{ continuous } \mathit{II}\text{-convex and} \\ \forall (p^*, q^*) \in P^* \times Q^*,\ w(p^*, q^*) \leq \mathrm{cav}_I \mathrm{Min}(w, \hat{v})(p^*, q^*) \end{matrix} \right\},$$

and a similar property for $\overline{v}^*$. The equality $\underline{v}^* = \overline{v}^*$ follows from Theorem 2.1 in [14] and implies $\overline{v} = \underline{v}$. □

Remark 6.3. *The above proof relies heavily on the fact that B is associated to a recurrent Markov chain. For example, recall example E of section 4:*

$$M = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 1/2 & 1/4 & 1/4 \end{pmatrix},\quad B = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 2/3 & 1/3 & 0 \end{pmatrix}.$$

In this case, the function $f(p) = (p^2 - 1/3)^+$ (where $^+$ denotes the positive part) is such that $\mathrm{cav} f(p_0) = 0$ *and* $\mathrm{cav}(f^*)(p_0 B) = 2/9$ *for $p_0 = (0, 0, 1)$. This was already pointed out in Renault [20], where general Markov matrices were considered. Our reduction of the problem to the study of aperiodic recurrent Markov chains allows therefore for useful technical simplifications. For example, in case L is reduced to a single point, our characterization becomes* $v(p) = \mathrm{cav}(\hat{v})(p)$. *Using aperiodic but possibly not recurrent Markov chains, the characterization given in [20] reduces to* $v(p) = \mathrm{cav}(\hat{v})(pB)$ *where $\hat{v}$ and B are defined similarly. One checks easily that these two expressions coincide.*

Remark 6.4. *Note that one may add the condition that w is also I-concave in the supremum defining $\underline{v}$, because if w fulfills the required assumptions, then* $\mathrm{cav}_I w$ *also and* $\mathrm{cav}_I w \geq w$. *A similar result holds for $\overline{v}$.*

Lemma 6.5. *Let $w \in \mathcal{W}$ be such that $\forall (p, q) \in \Delta(K) \times \Delta(L)$, $w(p, q) \leq \mathrm{cav}_I \mathrm{Min}(w, \hat{v})(p, q)$.*

If $w(p, q) > \hat{v}(p, q)$, then there exists a convex combination $(\alpha_m, p_m)_{m=1,..,K}$ in $[0, 1] \times \Delta(K)$ such that

$$\sum_{m=1}^{K} \alpha_m = 1,\ p = \sum_{m=1}^{K} \alpha_m p_m,\ \forall m = 1, .., K,\ w(p_m, q) \leq \hat{v}(p_m, q),\ \text{and} \quad \sum_{m=1}^{K} \alpha_m w(p_m, q) \geq w(p, q).$$

Proof. Using Caratheodory's Theorem, there exists $(\alpha_m, p_m)_{m=1,..,K}$ in $[0,1] \times \Delta(K)$ such that

$$\sum_{m=1}^{K} \alpha_m = 1,\ p = \sum_{m=1}^{K} \alpha_m p_m, \sum_{m=1}^{K} \alpha_m \mathrm{Min}(w, \hat{v})(p_m, q) = \mathrm{cav}_I \mathrm{Min}(w, \hat{v})(p, q).$$

This implies that for all $m = 1, .., K$,

$$\mathrm{Min}(w, \hat{v})(p_m, q) = \mathrm{cav}_I \mathrm{Min}(w, \hat{v})(p_m, q) \geq w(p_m, q),$$

so that $\mathrm{Min}(w, \hat{v})(p_m, q) = w(p_m, q) \leq \hat{v}(p_m, q)$. To conclude, note that

$$\sum_{m=1}^{K} \alpha_m w(p_m, q) = \mathrm{cav}_I \mathrm{Min}(w, \hat{v})(p, q) \geq w(p, q).$$

□

Lemma 6.6. *Consider a convex combination $p = \sum_{s \in S} \alpha_s p_s$ in $\Delta(K)$. Consider for each $s \in S$ a non revealing strategy σ_s in $\hat{\Sigma}(p_s)$. There exists σ in $\Sigma(p)$ such that:*

$$\forall q \in \Delta(L), \forall \tau \in \mathcal{T},\ \ \mathbb{P}_{p,q,\sigma,\tau} = \sum_{s \in S} \alpha_s \mathbb{P}_{p_s,q,\sigma_s,\tau}.$$

As a consequence we have for all $T \in \mathbb{N}^*$, q in $\Delta(L)$ and $\tau \in \mathcal{T}$,

$$\mathbb{E}_{p,q,\sigma,\tau}[\sum_{t=0}^{T-1} \|\hat{q}_{t+1}(q,\tau) - \hat{q}_t(q,\tau)\|] = \sum_{s \in S} \alpha_s \mathbb{E}_{p_s,q,\sigma_s,\tau}[\sum_{t=0}^{T-1} \|\hat{q}_{t+1}(q,\tau) - \hat{q}_t(q,\tau)\|].$$

Moreover, for any function $w \in \mathcal{W}$ which is I-concave, II-convex we have

$$\forall t \geq 1,\ \mathbb{E}_{p,q,\sigma,\tau}[w(\hat{p}_t, \hat{q}_t)] \geq \sum_{s \in S} \alpha_s w(p_s B, qC).$$

Proof. Let us define σ as in Lemma 5.6 via the splitting procedure: observe the first state k_1 in K, then choose a variable s in S according to the probability $\alpha_s p_s^{k_1}/p^{k_1}$, and play according to σ_s. It defines a probability $\mathbb{P}$ on $\Omega \times S$ and using Kuhn's Theorem, there exists σ such that the induced probability on Ω is equal to $\mathbb{P}_{p,q,\sigma,\tau}$. Moreover, for any event A in Ω, we have $\mathbb{P}(A|s) = \mathbb{P}_{p_s,q,\sigma_s,\tau}(A)$ almost surely, which implies the equality $\mathbb{P}_{p,q,\sigma,\tau} = \sum_{s \in S} \alpha_s \mathbb{P}_{p_s,q,\sigma_s,\tau}$.

Recall that whenever h_t is compatible with (q, τ), $q_t(q,\tau)(h_t)$ does not depend on (p, σ). Since $\hat{q}_t(q,\tau) = q_t(q,\tau)B$, the preceding equality implies that

$$\mathbb{E}_{p,q,\sigma,\tau}[\sum_{t=0}^{T-1} \|\hat{q}_{t+1}(q,\tau) - \hat{q}_t(q,\tau)\|] = \sum_{s \in S} \alpha_s \mathbb{E}_{p_s,q,\sigma_s,\tau}[\sum_{t=0}^{T-1} \|\hat{q}_{t+1}(q,\tau) - \hat{q}_t(q,\tau)\|].$$

Let us now work with the probability $\mathbb{P}$. Since σ_s is non-revealing at p_s, we have for all stage t, s in S and history h_t:

$$(\mathbb{P}(k_{t+1} = k | h_t, s))_{k \in K} B = (\mathbb{P}_{p_s,q,\sigma_s,\tau}(k_{t+1} = k | h_t))_{k \in K} B = \hat{p}_t(p_s, \sigma_s)(h_t) = p_s B.$$

Denote by $\mathbb{E}$ the expectation under $\mathbb{P}$, and by $\tilde{s}$ the random variable with values in S of the first choice of Player 1. The above line implies:

$$\mathbb{E}[p_{\tilde{s}}B|h_t] = \mathbb{E}[(\mathbb{P}(k_{t+1} = k|h_t, \tilde{s}))_{k\in K}|h_t]B = p_t(p,\sigma)(h_t)B = \hat{p}_t(p,\sigma)(h_t),$$

Using that w is I-concave and Jensen's inequality, we deduce that

$$\mathbb{E}[w(p_{\tilde{s}}B, \hat{q}_t(q,\tau))|h_t] \leq \mathbb{E}[w(\hat{p}_t(p,\sigma), \hat{q}_t(q,\tau))|h_t].$$

Using again that $\hat{q}_t(q,\tau)$ does not depend on (p,σ), we obtain

$$\mathbb{E}[w(p_{\tilde{s}}B, \hat{q}_t(q,\tau))] = \mathbb{E}[\mathbb{E}[w(p_{\tilde{s}}B, \hat{q}_t(q,\tau))|\tilde{s}]] = \sum_{s\in S} \alpha_s \mathbb{E}_{p_s,q,\sigma_s,\tau}[w(p_sB, \hat{q}_t(q,\tau)].$$

To conclude, w being $I\!I$-convex, and using Jensen's inequality,

$$\forall s \in S,\ \mathbb{E}_{p_s,q,\sigma_s,\tau}[w(p_sB, \hat{q}_t(q,\tau)] \geq w(p_sB, qC).$$

□

Let us now turn to the proof of the main theorem, making formal the description given in the end of section 4.

Proof of Theorem 6.1.
(1) *Fixing norms.* Recall that if $x = (x_s)_{s\in S}$ is an element of an euclidean space $\mathbb{R}^S$, $\|x\| = \sum_{s\in S} |x_s|$. For $T \in \mathbb{N}^*$, the set of strategies Σ_T is the set of maps from $\cup_{t=0}^{T-1}(K \times I \times J)^{t-1} \times K$ to $\Delta(I)$ which is seen as a subset of $\mathbb{R}^I$. Σ_T can be seen as a compact subset of $(\mathbb{R}^I)^{\cup_{t=0}^{T-1}(K\times I\times J)^{t-1}\times K}$ endowed with the corresponding norm. Note at first that for all $T \in \mathbb{N}^*$, for all $(p,q) \in \Delta(K) \times \Delta(L)$, and for all $\sigma \in \Sigma_T$ and $\tau, \tau' \in \mathcal{T}_T$,

$$|\gamma_T^{p,q}(\sigma,\tau) - \gamma_T^{p,q}(\sigma,\tau')| \leq \|\tau - \tau'\|.$$

(2) *Here we show that we can approximate strategies with low variations of the martingales $\hat{p}_t$ and $\hat{q}_t$ by non revealing strategies.* Let T be fixed. For $\frac{1}{I} > \varepsilon > 0$, define Σ_T^ε as the set of strategies $\sigma \in \Sigma_T$ such that at any stage and for any history of the game, all the actions are played with probability at least ε. Formally,

$$\forall t = 1,..,T, \forall h_{t-1} \in H_{t-1}, \forall (k_1,...,k_{t-1},k) \in K^t, \forall i \in I, \sigma_t(h_{t-1},k_1,..,k_{t-1})^k(i) \geq \varepsilon.$$

Define the map F_ε from $R_\varepsilon = \Delta(K) \times \Delta(L) \times \Sigma_T^\varepsilon \times \mathcal{T}_T$ to $\mathbb{R}^{\vec{T}}$ by

$$F_\varepsilon(p,q,\sigma,\tau) = (\mathbb{P}_{p,q,\sigma,\tau}(h_t)(\hat{q}_{t+1}(q,\tau)(h_t) - \hat{q}_t(q,\tau)(h_{t-1})))_{t=0,..,T-1,\, h_t\in(I\times J)^t}\ ,$$

where for $h_t = (i_1, j_1, .., i_t, j_t)$, h_{t-1} is defined here as $(i_1, j_1, .., i_{t-1}, j_{t-1})$, and $\vec{T}$ is the large but finite integer $\vec{T} = L\sum_{t=0}^{T-1}(I \times J)^t$. Note that F is continuous and that R_ε is compact. Define the compact set $\hat{R}_\varepsilon = F_\varepsilon^{-1}(\{0\})$, it is the subset of R_ε consisting of elements (p,q,σ,τ) such that τ is non-revealing at q. This follows easily from the observation that if an history

h_t is compatible with (q,τ), then due to the definition of Σ^ε_T we have $I\!P_{p,q,\sigma,\tau}(h_t) > 0$ for all p and $\sigma \in \Sigma^\varepsilon_T$. Note also that by construction

$$\|F(p,q,\sigma,\tau)\| = \sum_{t=0}^{T-1} I\!E_{p,q,\sigma,\tau}[\|\hat{q}_{t+1}(q,\tau) - \hat{q}_t(q,\tau)\|].$$

Claim: For all $\varepsilon > 0$ and $T \in I\!N^*$, there exists $C_1 = C_1(T,\varepsilon)$ such that

$$\forall (p,q,\sigma,\tau) \in R_\varepsilon,\ \exists (\hat{p},\hat{q},\hat{\sigma},\hat{\tau}) \in \hat{R}_\varepsilon,\ \|(p,q,\sigma,\tau) - (\hat{p},\hat{q},\hat{\sigma},\hat{\tau})\| \leq \varepsilon + C_1\|F(p,q,\sigma,\tau)\|.$$

To prove this claim, assume by contradiction that for all $n \in I\!N^*$, there exists $(p_n,q_n,\sigma_n,\tau_n) \in R_\varepsilon$ such that for all $(\hat{p},\hat{q},\hat{\sigma},\hat{\tau}) \in \hat{R}_\varepsilon$, $\|(p_n,q_n,\sigma_n,\tau_n) - (\hat{p},\hat{q},\hat{\sigma},\hat{\tau})\| > \varepsilon + n\|F(p_n,q_n,\sigma_n,\tau_n)\|$. By compactness, we can extract a convergent subsequence with limit $(p^*,q^*,\sigma^*,\tau^*)$ such that $F(p^*,q^*,\sigma^*,\tau^*) = 0$ and $\|(p^*,q^*,\sigma^*,\tau^*) - (\hat{p},\hat{q},\hat{\sigma},\hat{\tau})\| \geq \varepsilon$. which is a contradiction.

(3) *Here we show that an optimal strategy of Player 1 in a non revealing game $\hat{\Gamma}_T(p,q)$ is good against strategies of Player 2 that are non revealing at q', where q' is close to q.*

Claim: for all $\varepsilon > 0$ and $T \in I\!N^*$ there exists $C_2 = C_2(T,\varepsilon)$ such that for all $(p,q) \in \Delta(K) \times \Delta(L)$, for all $\sigma \in \hat{\Sigma}_T(p)$ which is optimal in $\hat{\Gamma}_T(p,q)$, for for all $q' \in \Delta(L)$, and for all $\tau' \in \hat{\mathcal{T}}(q')$,

$$\gamma^{p,q}_T(\sigma,\tau') \geq -\varepsilon - C_2\|q - q'\| + \hat{v}_T(p,q).$$

For a proof, assume by contradiction that for all $n \in I\!N^*$, there exists $(p_n,q_n) \in \Delta(K) \times \Delta(L)$, $\sigma_n \in \hat{\Sigma}_T(p_n)$ which is optimal in $\hat{\Gamma}_T(p_n,q_n)$, $q'_n \in \Delta(L)$, and $\tau'_n \in \hat{\mathcal{T}}(q'_n)$ such that

$$\gamma^{p_n,q_n}_T(\sigma_n,\tau'_n) \leq -\varepsilon - n\|q_n - q'_n\| + \hat{v}_T(p_n,q_n).$$

By compactness, there exists a convergent subsequence with limit $(p^*,q^*,\sigma^*,q'^*,\tau'^*)$. Since the left hand-side of the above inequality is bounded below by -1, we have $q^* = q'^*$. Moreover, the sets of non-revealing strategies being closed, σ^* is non-revealing at p^* and τ^* is non-revealing at q^*. Using step (1), we have for all n,

$$\hat{v}_T(p_n,q_n) - \|\tau^* - \tau_n\| \leq \gamma^{p_n,q_n}_T(\sigma_n,\tau^*) - \|\tau^* - \tau_n\| \leq \gamma^{p_n,q_n}_T(\sigma_n,\tau'_n).$$

It follows that

$$\hat{v}_T(p^*,q^*) \leq -\varepsilon + \hat{v}_T(p^*,q^*),$$

a contradiction.

(4) *Here we control the error while perturbing an optimal strategy in $\hat{\Gamma}(p,q)$ by a non revealing completely mixed strategy.*

Let σ^*_T denote the strategy in Σ_T which plays an uniform distribution over I at all stages independently of the history of the game, notice that σ^*_T is non-revealing at all points in $\Delta(K)$. For all $(p,q) \in \Delta(K) \times \Delta(L)$, let $\sigma_T(p,q)$ denote an optimal strategy of Player 1 in $\hat{\Gamma}(p,q)$. Denote by $\sigma_{T,\varepsilon}(p,q)$ the strategy which plays: σ^*_T with probability ε, and $\sigma_T(p,q)$ with probability $1-\varepsilon$. The strategy $\sigma_{T,\varepsilon}(p,q)$ is non revealing at p, and we have for all $\tau \in \mathcal{T}_T$,

$$I\!P_{p,q,\sigma_{T,\varepsilon},\tau} = \varepsilon I\!P_{p,q,\sigma^*_T,\tau} + (1-\varepsilon) I\!P_{p,q,\sigma_T(p,q),\tau}.$$

Using step (2), for all $(p,q) \in \Delta(K) \times \Delta(L)$, and all $\tau \in \mathcal{T}_T$,

$$\exists (\hat{p}, \hat{q}, \hat{\sigma}, \hat{\tau}) \in \hat{R}_\varepsilon,\ \|(p,q,\sigma_{T,\varepsilon}(p,q),\tau) - (\hat{p},\hat{q},\hat{\sigma},\hat{\tau})\| \leq \varepsilon + C_2(T,\varepsilon)\|F(p,q,\sigma_{T,\varepsilon}(p,q),\tau)\|.$$

Using then (1) and (3),

$$\begin{aligned}
\gamma_T^{p,q}(\sigma_{T,\varepsilon}(p,q),\tau) &= \varepsilon \gamma_T^{p,q}(\sigma_T^*,\tau) + (1-\varepsilon)\gamma_T^{p,q}(\sigma_T(p,q),\tau) \\
&\geq -2\varepsilon - \|\tau - \hat{\tau}\| + \gamma_T^{p,q}(\sigma_T(p,q),\hat{\tau}) \\
&\geq -2\varepsilon - \|\tau - \hat{\tau}\| - \varepsilon - C_2(T,\varepsilon)\|q - \hat{q}\| + \hat{v}_T(p,q) \\
&\geq \hat{v}_T(p,q) - 3\varepsilon - (1 + C_2(T,\varepsilon))C_1(T,\varepsilon)\|F(p,q,\sigma_{T,\varepsilon}(p,q),\tau)\|.
\end{aligned}$$

(5) *Here we prove that:* $\text{liminf}_T v_T \geq \underline{v}$.

Let $w \in \mathcal{W}$ be a balanced I-concave, II-convex function such that

$$\forall (p,q) \in \Delta(K) \times \Delta(L),\ w(p,q) \leq \text{cav}_I \text{Min}(w,\hat{v})(p,q).$$

Fix $(p,q) \in \Delta(K) \times \Delta(L)$, we have to show that $\text{liminf}_T v_T(p,q) \geq w(p,q)$.

Let $\varepsilon > 0$ and choose $T_0 \in \mathbb{N}^*$ such that $\|\hat{v}_{T_0} - \hat{v}\|_\infty \leq \varepsilon$. For simplicity, we will write p_t for $p_t(p,\sigma)$, q_t for $q_t(q,\tau)$ and C_1, C_2 for $C_1(T_0,\varepsilon), C_2(T_0\varepsilon)$. Given $N \in \mathbb{N}^*$ and an optimal strategy $\tau \in \overline{\mathcal{T}}$ of Player 2 in $\Gamma_{NT_0}(p,q)$, we define a strategy σ as follows.

The set of stages is divided into consecutive blocks of length T_0. For $n \geq 0$, at the beginning of block n, i.e. at the beginning of stage $nT_0 + 1$, Player 1 determines his strategy for the block according to the random variables p_{nT_0} and q_{nT_0}. Two cases may occur.

If $\hat{v}(p_{nT_0}, q_{nT_0}) \geq w(p_{nT_0}, q_{nT_0})$, then σ play during the next T_0 stages the non revealing strategy $\sigma_{T_0,\varepsilon}(p_{nT_0}, q_{nT_0})$.

If $\hat{v}(p_{nT_0}, q_{nT_0}) < w(p_{nT_0}, q_{nT_0})$, using Lemma 6.5, there exists a convex combination $(\alpha_{n,m}, p_{n,m})_{m=1,..,K}$ in $[0,1] \times \Delta(K)$ which depends on (p_{nT_0}, q_{nT_0}), such that

$$\sum_{m=1}^{K} \alpha_{n,m} = 1,\ p_{nT_0} = \sum_{m=1}^{K} \alpha_{n,m} p_{n,m}, \forall m = 1,..,K,\ w(p_{n,m}, q_{nT_0}) \leq \hat{v}(p_{n,m}, q_{nT_0})$$

$$\text{and} \quad \sum_{m=1}^{K} \alpha_{n,m} w(p_{n,m}, q_{nT_0}) \geq w(p_{nT_0}, q_{nT_0}).$$

In this case Player 1 plays at block n the strategy $\overline{\sigma}_{T_0,\varepsilon}(p_{nT_0}, q_{nT_0}) \in \Sigma_{T_0}$ given by Lemma 6.6 and such that for all $\tau' \in \mathcal{T}_{T_0}$,

$$\mathbb{P}_{p_{nT_0}, q_{nT_0}, \overline{\sigma}_{T_0,\varepsilon}(p_{nT_0}, q_{nT_0}), \tau'} = \sum_{m=1}^{K+1} \alpha_{n,m} \mathbb{P}_{p_{n,m}, q_{nT_0}, \sigma_{T,\varepsilon}(p_{n,m}, q_{nT_0}), \tau'}.$$

This ends the definition of σ.

We have to show that the payoff $\gamma_{NT_0}(\sigma,\tau)$ is large. It can be written as

$$\begin{aligned}\gamma_{NT_0}(\sigma,\tau) &= \frac{1}{NT_0}\sum_{n=0}^{N-1}\mathbb{E}_{p,q,\sigma,\tau}[\sum_{t=nT_0+1}^{(n+1)T_0}\mathbb{E}_{p,q,\sigma,\tau}[g(k_t,l_t,i_t,j_t)|h_{nT_0}]]\\ &= \frac{1}{N}\sum_{n=0}^{N-1}\mathbb{E}_{p,q,\sigma,\tau}[\gamma_{T_0}^{p_{nT_0},q_{nT_0}}(\sigma(h_{nT_0}),\tau(h_{nT_0}))]\end{aligned}$$

where $\tau(h_{nT_0})\in\mathcal{T}_{T_0}$ and $\sigma(h_{nT_0})\in\Sigma_{T_0}$ are the continuation strategies defined in Definition 5.14.

Claim: For all $n=0,..,N-1$ and h_{nT_0} occurring with positive probability,

$\gamma_{T_0}^{p_{nT_0},q_{nT_0}}(\sigma(h_{nT_0}),\tau(h_{nT_0}))\geq w(p_{nT_0},q_{nT_0})-3\varepsilon-(1+C_2)C_1\|F(p_{nT_0},q_{nT_0},\sigma(h_{nT_0}),\tau(h_{nT_0})\|.$

Proof of the claim. For the case $\hat{v}(p_{nT_0},q_{nT_0})\geq w(p_{nT_0},q_{nT_0})$, we have $\sigma(h_{nT_0})=\sigma_{T_0,\varepsilon}(p_{nT_0},q_{nT_0})$ and

$$\begin{aligned}\gamma_{T_0}^{p_{nT_0},q_{nT_0}}(\sigma_{T_0,\varepsilon}(p_{nT_0},q_{nT_0}),\tau(h_{nT_0}))&\geq\hat{v}_{T_0}(p_{nT_0},q_{nT_0})-3\varepsilon-(1+C_2)C_1\|F(p_{nT_0},q_{nT_0},\sigma(h_{nT_0}),\tau(h_{nT_0})\|\\ &\geq w(p_{nT_0},q_{nT_0})-3\varepsilon-(1+C_2)C_1\|F(p_{nT_0},q_{nT_0},\sigma(h_{nT_0}),\tau(h_{nT_0})\|.\end{aligned}$$

For the case $\hat{v}(p_{nT_0},q_{nT_0})<w(p_{nT_0},q_{nT_0})$, we have $\sigma(h_{nT_0})=\overline{\sigma}_{T_0,\varepsilon}(p_{nT_0},q_{nT_0})$ and

$$\begin{aligned}\gamma_{T_0}^{p_{nT_0},q_{nT_0}}(\overline{\sigma}_{T_0,\varepsilon}(p_{nT_0},q_{nT_0}),\tau(h_{nT_0}))&=\sum_{m=1}^{K}\alpha_{n,m}\gamma_{T_0}^{p_{nT_0},q_{nT_0}}(\sigma_{T,\varepsilon}(p_{n,m},q_{nT_0}),\tau(h_{nT_0}))\\ &\geq\sum_{m=1}^{K}\alpha_{n,m}\hat{v}_{T_0}(p_{n,m},q_{nT_0})-3\varepsilon-(1+C_2)C_1\sum_{m=1}^{K}\alpha_{n,m}\|F(p_{n,m},q_{nT_0},\sigma_{T,\varepsilon}(p_{n,m},q_{nT_0}),\tau(h_{nT_0})\|\\ &\geq w((p_{nT_0},q_{nT_0})-3\varepsilon-(1+C_2)C_1\|F(p_{nT_0},q_{nT_0},\overline{\sigma}_{T,\varepsilon}(p_{nT_0},q_{nT_0})),\tau(h_{nT_0})\|,\end{aligned}$$

where the last inequality follows from the Lemmas 6.5 and 6.6. □

Let us now prove that:

$$\forall n=1,..,N-1,\quad\mathbb{E}_{p,q,\sigma,\tau}[w(p_{nT_0},q_{nT_0})]\geq w(p,q).\tag{4}$$

By induction, it is sufficient to prove that for $n=0,..,N-2$,

$$\mathbb{E}_{p,q,\sigma,\tau}[w(p_{(n+1)T_0},q_{(n+1)T_0})|h_{nT_0}]\geq w(p_{nT_0},q_{nT_0}).$$

Let us at first consider the case $\hat{v}(p_{nT_0},q_{nT_0})\geq w(p_{nT_0},q_{nT_0})$. Since $\sigma(h_{nT_0})$ is non-revealing at p_{nT_0}, we have $p_{(n+1)T_0}=p_{nT_0}$. Then, using that w is balanced, it is sufficient to prove that

$$\mathbb{E}_{p,q,\sigma,\tau}[w(p_{nT_0}B,q_{(n+1)T_0}B)|h_{nT_0}]\geq w(p_{nT_0}B,q_{nT_0}B).$$

The process $\hat{q}_t = q_t B$ being a martingale, the above inequality follows from the fact that w is $I\!I$-convex and Jensen's inequality. Let us now consider the case $\hat{v}(p_{nT_0}, q_{nT_0}) < w(p_{nT_0}, q_{nT_0})$. Using Lemma 6.6, we have

$$\begin{aligned}
&I\!E_{p,q,\sigma,\tau}[w(p_{(n+1)T_0}B, q_{(n+1)T_0}B)|h_{nT_0}] \\
&\qquad = I\!E_{p_{nT_0}, q_{nT_0}, \overline{\sigma}_{T,\varepsilon}(p_{nT_0}, q_{nT_0}), \tau(h_{nT_0})}[w(\hat{p}_{T_0}(p_{nT_0}, \overline{\sigma}_{T,\varepsilon}(p_{nT_0}, q_{nT_0})), \hat{q}_{T_0}(q_{nT_0}, \tau(h_{nT_0}))] \\
&\qquad \geq \sum_{m=1}^{K} \alpha_{n,m} w(p_{n,m}B, q_{nT_0}B) = \sum_{m=1}^{K} \alpha_{n,m} w(p_{n,m}, q_{nT_0}) \geq w(p_{nT_0}, q_{nT_0}).
\end{aligned}$$

This proves the above inequalities (4).

Summing up, we have:

$$\gamma_{NT_0}(\sigma,\tau) \geq w(p,q) - 3\varepsilon - (1+C_2)C_1 \frac{1}{N} \sum_{n=0}^{N-1} I\!E_{p,q,\sigma,\tau}[\|F(p_{nT_0}, q_{nT_0}, \sigma(h_{nT_0}), \tau(h_{nT_0})\|.$$

Finally, using Lemma 5.15, and the classical bound on the L_1-variation of martingales (see for instance Proposition 3.8. in [29]), we obtain:

$$\begin{aligned}
\sum_{n=0}^{N-1} I\!E_{p,q,\sigma,\tau}[\|F(p_{nT_0}, q_{nT_0}, \sigma(h_{nT_0}), \tau(h_{nT_0})\|] &= \sum_{n=0}^{N-1} I\!E_{p,q,\sigma,\tau}[I\!E_{p,q,\sigma,\tau}[\sum_{t=nT_0+1}^{(n+1)T_0} \|\hat{q}_t - \hat{q}_{t-1}\| | h_{nT_0}]] \\
&= \sum_{t=1}^{NT_0} I\!E_{p,q,\sigma,\tau}[\|\hat{q}_t - \hat{q}_{t-1}\|] \\
&\leq \sqrt{NT_0(L-1)}.
\end{aligned}$$

We conclude that:

$$\gamma_{NT_0}(\sigma,\tau) \geq w(p,q) - 3\varepsilon - (1+C_2)C_1 \frac{\sqrt{T_0(L-1)}}{\sqrt{N}}.$$

Since τ is optimal in $\Gamma_{NT_0}(p,q)$, we deduce that

$$v_{NT_0}(p,q) \geq w(p,q) - 3\varepsilon - (1+C_2)C_1 \frac{\sqrt{T_0(L-1)}}{\sqrt{N}}.$$

and then

$$\text{liminf}_N v_{NT_0}(p,q) \geq w(p,q) - 3\varepsilon.$$

Using that $\|v_T - v_{T+T'}\|_\infty \leq \frac{2T'}{T}$, this implies

$$\text{liminf}_T v_T(p,q) \geq w(p,q) - 3\varepsilon,$$

and finally $\text{liminf}_T v_T(p,q) \geq w(p,q)$. We deduce that $\text{liminf}_T v_T(p,q) \geq \underline{v}$.

(6) *We conclude the proof of Theorem 6.1.* Symmetrically, we obtain as in point (5) that $\text{limsup}_T v_T(p,q) \leq \overline{v}$. The conclusion follows therefore from Proposition 6.2. ☐

7 Proof of Proposition 5.13

Let us begin with a few definitions. Recall that K is partitioned into recurrence classes $K(1)$, ..., $K(r_M)$, each recurrent class r having a unique invariant measure $p^*(r) \in \Delta(K(r))$. $P^* = \text{conv}\{p^*(1), ..., p^*(M)\}$ is the set of invariant measures for M, as well as for $B = \lim_t M^t$. In the following we denote by $\Delta(r_M)$ the set of probabilities over the recurrences classes of M, i.e. over the finite set $\{1, ..., r_M\}$.

Notation 7.1. *Let p be in $\Delta(K)$.*

For each r, we denote by $\lambda(p)^r = \sum_{k \in K(r)} p^k$ the probability of the recurrence class r under p. And we define the probability $\lambda(p) \in \Delta(r_M)$ by $\lambda(p) = (\lambda(p)^r)_{r=1,...,r_M}$.

For each $r = 1, .., r_M$, we denote the conditional law, under p, of k given r as the vector $p_{|r} \in \Delta(K)$ such that:

$$p^k_{|r} = \begin{cases} \frac{p^k}{\lambda(p)^r} & \text{if} \quad \lambda(p)^r > 0\,,\ k \in K(r), \\ p^*(r)^k & \text{if} \quad \lambda(p)^r = 0,\ k \in K(r), \\ 0 & \text{if} \qquad\quad k \notin K(r). \end{cases}$$

It follows that for all $p \in \Delta(K)$, $p = \sum_{r=1}^{r_M} \lambda(p)^r p_{|r}$. Since $p^k_{|r} = 0$ for $k \notin K(r)$, $p_{|r}$ will be often assimilated to a vector in $\Delta(K(r))$.

Definition 7.2. *The map $S : \Delta(K) \times \Delta(K) \to I\!\!R_+$ is defined by:*

$$\forall p, p' \in \Delta(K), \ \ S(p, p') = \|\lambda(p) - \lambda(p')\| + \sum_{r\,:\,\lambda(p)^r \lambda(p')^r > 0} \lambda(p')^r \ \|p_{|r} - p'_{|r}\|.$$

The asymmetric map S will play an important role in the sequel, and will be used as a (quasi, semi-) metric. S is clearly not a metric, but separates points (we have $S(p, p') = 0$ if and only if $p = p'$) and enjoys the following useful properties.

Lemma 7.3. *For all $p, p' \in \Delta(K)$,*

$$\|p - p'\| \leq S(p, p') \leq 3\|p - p'\|, \ S(pM, p'M) \leq S(p, p'),$$

$$S(pB, p'B) = \|pB - p'B\| = \|\lambda(p) - \lambda(p')\|, \ \text{and} \ \ S(p, p') = \|p - p'\| \ \text{if} \ \ pB = p'B.$$

Before proving the lemma, we will present the roadmap of the proof

Definition 7.4.

1) For $\alpha \in [0, 1]$, define the operator Φ_α on the set $\mathcal{W}$ of continuous functions f defined on $\Delta(K) \times \Delta(L)$ by

$$\Phi_\alpha(f)(p, q) = \max_{x \in NR(p)} \min_{y \in NR(q)} \left(\alpha\, G(p, q, x, y) + (1 - \alpha) \sum_{i \in I, j \in J} x(p)(i) y(q)(j) f(\vec{p}(x, i)M, \vec{q}(y, j)N) \right).$$

2) Let $\mathcal{F}$ be the subset of functions f in $\mathcal{W}$ satisfying:

- $\forall p, p' \in \Delta(K)$, $\forall q \in \Delta(L)$, $f(p,q) - f(p',q) \leq S(p,p')$,
- $\forall p \in \Delta(K)$, $\forall q, q' \in \Delta(L)$, $f(p,q) - f(p,q') \leq S(q',q)$,
- $\forall p^*, q^* \in P^* \times Q^*$, f *is I-concave and II-convex on* $B^{-1}(p^*) \times C^{-1}(q^*)$.

We now present our strategy for the proof of Proposition 5.13. We have $\hat{v}_{t+1} = \Phi_{\frac{1}{t+1}}(\hat{v}_t)$ for $t \geq 0$ (with the convention $\hat{v}_0 = 0$), and all the functions in $\mathcal{F}$ are 3-Lipschitz using Lemma 7.3. As a consequence, to show that $\hat{v}_t$ is 3-Lipschitz for each t, it is sufficient to show that for all α, $\mathcal{F}$ is stable by Φ_α.

To do so, the main point is the following: given p and p' in $\Delta(K)$ and x in $NR(p)$, one need to find p' in $NR(p')$ such that (p', x') is not too far from (p, x). If p and p' belong to the same set $B^{-1}(p^*)$, i.e. if p and p' assigns the same weight to each recurrence class, this can be efficiently done using, within each simplex $\Delta(K(r))$, a lemma by Laraki ([11], see here the Appendix at the end). If $pB \neq p'B$, there exists a simple affine map to transfer probabilities in $B^{-1}(pB)$ to probabilities in $B^{-1}(p'B)$ (Lemma 7.12). In the general case, we need to combine both aspects, applying first the affine transformation then the Laraki splitting (see Lemma 7.13). In the computations, the expression $S(p,p')$ appears and allows to control the expressions.

We now prove Lemma 7.3. After this, all the rest of this section is devoted to the proof that:

$$\forall \alpha \in [0,1], \quad \Phi_\alpha(\mathcal{F}) \subset \mathcal{F}.$$

Proof of Lemma 7.3.

$$\begin{aligned}
\|p - p'\| &= \sum_{r\,:\,\lambda(p)^r\lambda(p')^r>0} \; \sum_{k\in K(r)} |\lambda(p)^r p^k_{|r} - \lambda(p')^r p'^k_{|r}| \\
&\quad + \sum_{r\,:\,\lambda(p)^r=0\,,\,\lambda(p')^r>0} \; \sum_{k\in K(r)} p'^k + \sum_{r\,:\,\lambda(p)^r>0\,,\,\lambda(p')^r=0} \; \sum_{k\in K(r)} p^k \\
&\leq \sum_{r\,:\,\lambda(p)^r\lambda(p')^r>0} \; \sum_{k\in K(r)} \Big(|\lambda(p)^r p^k_{|r} - \lambda(p')^r p^k_{|r}| + |\lambda(p')^r p^k_{|r} - \lambda(p')^r p'^k_{|r}| \Big) \\
&\quad + \sum_{r\,:\,\lambda(p)^r=0\,,\,\lambda(p')^r>0} \lambda(p')^r + \sum_{r\,:\,\lambda(p)^r>0\,,\,\lambda(p')^r=0} \lambda(p)^r \\
&= \sum_{r=1}^{r_M} |\lambda(p)^r - \lambda(p')^r| + \sum_{r\,:\;\lambda(p)^r\lambda(p')^r>0} \lambda(p')^r \sum_{k\in K(r)} |p^k_{|r} - p'^k_{|r}| \\
&= S(p,p').
\end{aligned}$$

On the other hand,

$$\begin{aligned}
S(p,p') &= \|\lambda(p)-\lambda(p')\| + \sum_{r\,:\,\lambda(p)^r\lambda(p')^r>0} \lambda(p')^r \|p_{|r} - p'_{|r}\| \\
&\leq \|\lambda(p)-\lambda(p')\| + \sum_{r\,:\,\lambda(p)^r\lambda(p')^r>0} \sum_{k\in K(r)} \Big(|\lambda(p')^r p^k_{|r} - p^k| + |p^k - p'^k| \Big) \\
&= \|\lambda(p)-\lambda(p')\| + \sum_{r\,:\;\lambda(p)^r\lambda(p')^r>0} \left(|\lambda(p')^r - \lambda(p)^r| + \sum_{k\in K(r)} |p^k - p'^k| \right) \\
&\leq 2\|\lambda(p)-\lambda(p')\| + \|p-p'\| \leq 3\|p-p'\|.
\end{aligned}$$

For the next inequality, note that $\lambda(pM) = \lambda(p)$ and $(pM)_{|r} = p_{|r}M$, so that using that $p \to pM$ is non-expansive

$$\begin{aligned}
S(pM,p'M) &= \|\lambda(p)-\lambda(p')\| + \sum_{r\,:\,\lambda(p)^r\lambda(p')^r>0} \lambda(p')\|(p_{|r} - p'_{|r})M\| \\
&\leq \|\lambda(p)-\lambda(p')\| + \sum_{r\,:\,\lambda(p)^r\lambda(p')^r>0} \lambda(p')\|(p_{|r} - p'_{|r})\| = S(p,p').
\end{aligned}$$

The two last equalities follow easily by a direct computation. □

Fix p in $\Delta(K)$ and q in $\Delta(L)$, and let $p^* = pB$ and $q^* = qC$. To any $x \in \Delta(I)^K$, one associates $z \in \Delta(K \times I)$ by $z(k,i) = p^k x^k(i)$ so that $\vec{p}(z,i) = \vec{p}(x,i)$. We will need a more abstract way to denote conditional probabilities. Let $\Delta_f(\Delta(K))$ denote the set of probabilities μ with finite support on $\Delta(K)$. Such a probability can be written as

$$\mu = \sum_{n=1}^{N} \alpha_n \delta_{p_n},$$

with $(\alpha_n, p_n)_{n=1,..,N} \in ([0,1] \times \Delta(K))^N$ such that $\sum_{n=1}^N \alpha_n = 1$ and $p = \sum_{n=1,..,N} \alpha_n p_n$. This set is endowed with the usual weak* topology which is in particular metrized by the Wasserstein (Kantorovich-Rubinstein) distance induced by the norm $\|.\|$ and denoted d_W. Recall that

$$\forall \mu,\nu \in \Delta(_f(\Delta(K)),\ d_W(\mu,\nu) = \max\{\int \phi d\mu - \int \phi d\nu \mid \phi \in Lip_1\},$$

where Lip_1 denotes the set of 1-Lipschitz functions on $\Delta(K)$. We will use this particular metric on $\Delta_f(\Delta(K))$. Let us also recall the classical duality theorem (see e.g. Theorem 1.14 in [28]) (*Kantorovich duality formula*):

$$\forall \mu,\nu \in \Delta_f(\Delta(K)),\ d_W(\mu,\nu) = \min\{\int_{p,p'\in\Delta(K)} \|p-p'\| d\pi(p,p') \mid \pi \in P(\mu,\nu)\}$$

where $P(\mu,\nu)$ denotes the set of probability distributions on $\Delta(K) \times \Delta(K)$ having μ,ν for marginals.

Definition 7.5. *Let* $\Psi : \Delta(K \times I) \to \Delta_f(\Delta(K))$ *be defined by*

$$\Psi(z) = \sum_{i \in I\,:\, z(i)>0} z(i)\delta_{\vec{p}(z,i)}$$

with $z(i) = \sum_{k \in K} z(k,i)$ *and* $\vec{p}(z,i)^k = \frac{z(k,i)}{z(i)}$ *for* i *such that* $z(i) > 0$.

Ψ is a disintegration mapping. Imagine (k,i) is selected according to z and only i is observed, then for each i one can compute the conditional $\vec{p}(z,i)$ on K. $\Psi(z)$ gives the law of this posterior on K. Using the z variable and the desintegration Ψ, the definition of $NR(p)$ can now be read as follows:

$$x \in NR(p) \Longleftrightarrow \forall k \in K, \sum_{i \in I} z(k,i) = p^k \text{ and } \Psi(z) \in \Delta_f(B^{-1}(p^*)).$$

Let us now recall a few properties of Ψ.

Definition 7.6. $\Delta_f(\Delta(K))$ *is endowed with the convex order denoted* $\leq$ *and defined by*

$$\mu \leq \nu \Longleftrightarrow \int \phi d\mu \leq \int \phi d\nu \text{ for all l.s.c. convex } \phi : \Delta(K) \to I\!R \cup \{+\infty\}.$$

For instance, given p and p' in $\Delta(K)$ the Dirac measure $\delta_{(p+p')/2}$ is smaller than the average $1/2\ \delta_p + 1/2\ \delta_{p'}$.

Lemma 7.7. Ψ *is continuous and convex.*

Proof. See e.g. [21] Lemmas 4.16 and 4.17. □

The convexity of Ψ with respect to the convex order can be interpreted as the positive value of having more information. Before going back to the main proof, let us also recall the classical relationship between convex order and martingales.

Theorem 7.8 (Blackwell,1953 [2])**.** *For all* $\mu, \nu \in \Delta_f(\Delta(K))$, *if* U *and* V *denote respectively the supports of* μ *and* ν, *then*

$$\mu \leq \nu \Leftrightarrow \begin{cases} \exists F : U \to \Delta(V) \text{ s.t. :} \\ \forall u \in U,\ \sum_{v \in V} vF(u)(v) = u, \\ \sum_{u \in U} \mu(u)F(u) = \nu. \end{cases}$$

The condition on the right hand side can also be seen as follows: μ is the law of some random variable X_1 with values in $\Delta(K)$, ν is the law of some random variable X_2 with values in $\Delta(K)$, and we have the martingale condition: $I\!E(X_2|X_1) = X_1$.

We will make a change of variable in the expression of $\Phi_\alpha(f)$. The definition of $NR(p)$ can now be stated as:

$$x \in NR(p) \Longleftrightarrow \delta_p \leq \Psi(z) \text{ and } \Psi(z) \in \Delta_f(B^{-1}(p^*)).$$

Definition 7.9. *We define the set-valued maps:*

$$\begin{array}{rrcl} H: & \Delta(K) & \rightrightarrows & \Delta_f(\Delta(K)) \\ & p \in \Delta(K) & \to & H(p) = \{\mu \in \Delta_f(B^{-1}(pB)) | \int_{\Delta(K)} p' d\mu(p') = p\}, \end{array}$$

$$\begin{array}{rrcl} R: & \Delta_f(\Delta(K)) & \rightrightarrows & \Delta(K \times I) \\ & \mu \in \Delta_f(\Delta(K)) & \to & R(\mu) = \{z \in \Delta(K \times I) \,:\, \Psi(z) \leq \mu\}. \end{array}$$

We define similarly $H : \Delta(L) \rightrightarrows \Delta_f(\Delta(L))$ *and* $R : \Delta_f(\Delta(L) \rightrightarrows \Delta(L \times J)$ *(keeping the same letters* H *and* R *for simplicity).*

$H(p)$ is the set of probabilities μ over $B^{-1}(pB)$ with mean p, equivalently it is the set of probabilities μ over $B^{-1}(pB)$ such that $\delta_p \leq \mu$. Hence $H(p)$ is the set of splittings at p such that all posterior keep the same weight on each recurrence class. The set $R(\mu)$ can be seen the set of "strategies" z with disintegration no more informative than μ.

Lemma 7.10. *For* $\mu \in H(p)$ *and* $z \in R(\mu)$*, we have* $\Psi(z) \in H(p)$*.*

Proof. Write $p^* = pB$. Consider the map ξ equal to 0 on $B^{-1}(p^*)$ and to $+\infty$ elsewhere. It is convex and l.s.c. since $B^{-1}(p^*)$ is closed and convex, and

$$0 \leq \int \xi d\Psi(z) \leq \int \xi d\mu = 0,$$

which implies $\Psi(z) \in \Delta_f(B^{-1}(p^*))$. Moreover, for all $k \in K$, the map $\tilde{p} \to \tilde{p}^k$ being also convex and l.s.c.,

$$\int \tilde{p}^k d\Psi(z)(\tilde{p}) \leq \int \tilde{p}^k d\mu(\tilde{p}) = p^k,$$

which implies $\int \tilde{p} d\Psi(z)(\tilde{p}) = p$ since it belongs to $\Delta(K)$. ☐

It follows from the above definitions and the characterization of $NR(p)$ that for all $\alpha \in [0,1]$ and f, one can rewrite $\Phi_\alpha(f)$ as follows:

$$\begin{aligned} &\Phi_\alpha(f)(p,q) \\ &\quad = \sup_{z \text{ s.t.} \Psi(z) \in H(p)} \inf_{w \text{ s.t.} \Psi(w) \in H(q)} \left(\alpha G(z,w) + (1-\alpha) \int_{\Delta(K) \times \Delta(L)} f(\tilde{p}M, \tilde{q}N) d\Psi(z) \otimes \Psi(w)(\tilde{p}, \tilde{q}) \right) \\ &\quad = \sup_{\mu \in H(p)} \sup_{z \in R(\mu)} \inf_{\nu \in H(q)} \inf_{w \in R(\nu)} \left(\alpha G(z,w) + (1-\alpha) \int_{\Delta(K) \times \Delta(L)} f(\tilde{p}M, \tilde{q}N) d\Psi(z) \otimes \Psi(w)(\tilde{p}, \tilde{q}) \right), \end{aligned}$$

where $G(z,w) = \sum_{(k,l,i,j) \in K \times L \times I \times J} z(k,i) w(l,j) g(k,l,i,j)$.

We will simplify the above expression as follows.

Proposition 7.11. *Define* $V(\mu,\nu) = \sup_{z \in R(\mu)} \inf_{w \in R(\nu)} G(z,w)$*. Then* V *is* 1*-Lipschitz for* d_W *and is I-concave and II-convex on* $\Delta_f(\Delta(K)) \times \Delta_f(\Delta(L))$*. Moreover:*

$$\Phi_\alpha(f)(p,q) = \sup_{\mu \in H(p)} \inf_{\nu \in H(q)} \left(\alpha V(\mu,\nu) + (1-\alpha) \int_{\Delta(K) \times \Delta(L)} f(\tilde{p}M, \tilde{q}N) d\mu \otimes \nu(\tilde{p}, \tilde{q}) \right).$$

Proof of Proposition 7.11: Using that f is *II*-convex on $B^{-1}(p^*) \times C^{-1}(q^*)$ and that $C^{-1}(q^*)$ is invariant by N, we have for all $w \in R(\nu)$

$$\int_{\Delta(K\times\Delta(L)} f(\tilde{p}M, \tilde{q}N) d\Psi(z) \otimes \Psi(w)(\tilde{p}, \tilde{q}) \leq \int_{\Delta(K)\times\Delta(L)} f(\tilde{p}M, \tilde{q}N) d\Psi(z) \otimes \nu(\tilde{p}, \tilde{q}),$$

which implies the following equality

$$\inf_{\nu\in H(q)} \inf_{w\in R(\nu)} \left(\alpha G(z,w) + (1-\alpha)\int_{\Delta(K)\times\Delta(L)} f(\tilde{p}M, \tilde{q}N) d\Psi(z) \otimes \Psi(w)(\tilde{p}, \tilde{q})\right)$$
$$= \inf_{\nu\in H(q)} \inf_{w\in R(\nu)} \left(\alpha G(z,w) + (1-\alpha)\int_{\Delta(K)\times\Delta(L)} f(\tilde{p}M, \tilde{q}N) d\Psi(z) \otimes \nu(\tilde{p}, \tilde{q})\right)$$

since one can always choose $\nu = \Psi(w)$ in the right-hand side. A similar equality holds for $\Psi(z)$ and μ, and this implies that for the moment we have proved:

For all $\alpha \in [0,1]$, f continuous and $(p,q) \in \Delta(K)$,

$$\Phi_\alpha(f)(p,q) = \sup_{\mu\in H(p)} \sup_{z\in R(\mu)} \inf_{\nu\in H(q)} \inf_{w\in R(\nu)} \left(\alpha G(z,w) + (1-\alpha)\int_{\Delta(K)\times\Delta(L)} f(\tilde{p}M, \tilde{q}N) d\mu \otimes \nu(\tilde{p}, \tilde{q})\right).$$

Let us now study the properties of the set-valued map R.

Claim. The set-valued map $R : \mu \to R(\mu)$ is non expansive from $(\Delta_f(\Delta(K)), d_W)$ to $(\Delta(K \times I), \|.\|)$. Moreover, for all $\beta \in [0,1]$, and $\mu, \mu' \in \Delta_f(\Delta(K))$,

$$\beta R(\mu) + (1-\beta)R(\mu') \subset R(\beta\mu + (1-\beta)\mu').$$

Similar properties also hold for $R : \nu \to R(\nu)$ from $\Delta_f(\Delta(L))$ to $\Delta(L \times J)$.

Proof of the claim: We first prove the non expansive property. Fix μ and $\mu' \in \Delta_f(\Delta(K))$, and $z \in R(\mu)$, we have to find z' in $R(\mu')$ such that $\|z - z'\| \leq d_W(\mu, \mu')$. The idea is to link $\Psi(z)$ to μ via Theorem 7.8, then to link μ and μ' via Kantorovich duality formula, and finally to define z' using both links.

Since $\Psi(z) \leq \mu$, there exists a map $F : \Delta(K) \to \Delta_f(\Delta(K))$ such that for all $x \in \Delta(K)$, $F(x)$ is centered in x and $\mu = \sum_{i\in I} z(i) F(\vec{p}(z,i))$. Let U, V denote respectively the supports of μ and μ', and consider by Kantorovich duality a probability π on $U \times V$ with marginals (μ, μ') such that $\sum_{(u,v)\in U\times V} \|u - v\| \pi(u,v) = d_W(\mu, \mu')$. For $u \in U$, let $\vec{\pi}(u) = (\frac{\pi(u,v)}{\mu(u)})_{v\in V}$ denote the conditional law on V given u induced by π. Because the second marginal of π is μ', we have $\sum_{u\in U} \mu(u)\vec{\pi}(u) = \mu'$. We define, for all i in I:

$$\vec{p'}(i) = \sum_{v\in V} v\ m'(i)(v) \in \Delta(K) \text{ with } m'(i) = \sum_{u\in U} F(\vec{p}(z,i))(u)\ \vec{\pi}(u) \in \Delta(V),$$

and $z' \in \Delta(K \times I)$ by $z'(k,i) = z(i)\ \vec{p'}(i)^k \quad \forall k \in K, \forall i \in I$.

We have $\sum_{i\in I} z(i)m'(i) = \mu'$, and for each i the probability $m'(i)$ is centered in $\vec{p'}(i)$. By construction $\Psi(z') = \sum_{i\in I} z(i)\delta_{\vec{p'}(i)}$. All this implies by Theorem 7.8 that $\Psi(z') \leq \mu'$. Using Jensen's inequality, we have

$$\begin{aligned}
\|z - z'\| &= \sum_{i\in I} z(i)\|\vec{p}(z,i) - \vec{p'}(i)\| \\
&= \sum_{i\in I} z(i)\| \sum_{u\in U} uF(\vec{p}(z,i))(u) - \sum_{(u,v)\in U\times V} vF(\vec{p}(z,i))(u)\vec{\pi}(u)(v)\| \\
&= \sum_{i\in I} z(i)\| \sum_{u\in U}(u - \sum_{v\in V} v\vec{\pi}(u)(v))F(\vec{p}(z,i))(u)\| \\
&\leq \sum_{i\in I} z(i) \sum_{u\in U} F(\vec{p}(z,i))(u)\|u - \sum_{v\in V} v\vec{\pi}(u)(v)\| \\
&\leq \sum_{i\in I} z(i) \sum_{u\in U} F(\vec{p}(z,i))(u) \sum_{v\in V} \vec{\pi}(u)(v)\|u - v\| \\
&= \sum_{u\in U}\sum_{v\in V}(\sum_{i\in I} z(i)F(\vec{p}(z,i))(u))\vec{\pi}(u)(v)\|u - v\| \\
&= \sum_{u\in U}\sum_{v\in V} \mu(u)\vec{\pi}(u)(v)\|u - v\| \\
&= \sum_{(u,v)\in U\times V} \pi(u,v)\|u - v\| \\
&= d_W(\mu,\mu')
\end{aligned}$$

For the second assertion, let $z \in R(\mu)$ and $z' \in R(\mu')$, then since Ψ is convex,

$$\Psi(\beta z + (1-\beta)z') \leq \beta\Psi(z) + (1-\beta)\Psi(z') \leq \beta\mu + (1-\beta)\mu'.$$

This ends the proof of the claim, and we can conclude the proof of Proposition 7.11.

The law μ being fixed, let us now consider the map

$$(z,\nu) \in R(\mu) \times H(q) \to \inf_{w\in R(\nu)} \left(\alpha G(z,w) + (1-\alpha)\int_{\Delta(K)\times\Delta(L)} f(\tilde{p}M,\tilde{q}N)d\mu\otimes\nu(\tilde{p},\tilde{q}) \right).$$

As an infimum of linear continuous functions, it is concave and upper semi-continuous with respect to z on the compact convex set $R(\mu)$. On the other hand, the map

$$\nu \to \int_{\Delta(K)\times\Delta(L)} f(\tilde{p}M,\tilde{q}N)d\mu\otimes\nu(\tilde{p},\tilde{q})$$

is linear and using the second point in the above claim, the map

$$\nu \to \inf_{w\in R(\nu)} G(z,w)$$

is convex with respect to ν. Indeed, let $\beta \in [0,1]$ and $\nu,\nu' \in H(q)$. By compactness, there exist w^*, w'^* which minimize $G(z,.)$ on $R(\nu)$ and $R(\nu')$ respectively. Since $\beta w + (1-\beta)w' \in$

$R(\beta\nu + (1-\beta)\nu')$,

$$\begin{aligned}\beta \min_{w\in R(\nu)} G(z,w) + (1-\beta)\min_{w\in R(\nu')} G(z,w) &= \beta G(z,w^*) + (1-\beta)G(z,w'^*)\\ &= G(z,\beta w^* + (1-\beta)w'^*)\\ &\geq \min_{w\in R(\beta\nu+(1-\beta)\nu')} G(z,w).\end{aligned}$$

It allows to apply Sion's Minmax Theorem (see [24]) to conclude that

$$\Phi_\alpha(f)(p,q) = \sup_{\mu\in H(p)} \inf_{\nu\in H(q)} \sup_{z\in R(\mu)} \inf_{w\in R(\nu)} \left(\alpha G(z,w) + (1-\alpha)\int_{\Delta(K)\times\Delta(L)} f(\tilde{p}M,\tilde{q}N)d\mu\otimes\nu(\tilde{p},\tilde{q})\right).$$

Using the claim about R and that $z \to G(z,w)$ is 1-Lipschitz, the function V is 1-Lipschitz for d_W. Moreover, using the same proof as above, V is *I*-concave and *II*-convex on $\Delta_f(\Delta(K)) \times \Delta_f(\Delta(L))$. The preceding expression becomes

$$\Phi_\alpha(f)(p,q) = \sup_{\mu\in H(p)} \inf_{\nu\in H(q)} \left(\alpha V(\mu,\nu) + (1-\alpha)\int_{\Delta(K)\times\Delta(L)} f(\tilde{p}M,\tilde{q}N)d\mu\otimes\nu(\tilde{p},\tilde{q})\right),$$

and this concludes the proof of Proposition 7.11. □

It remains now to study the properties of the set-valued map H. Let us start with the following lemma which proves a kind of "triangle equality" for the map S.

Lemma 7.12. *For any pair (p^*, p'^*) in P^*, there exists an affine map $L = L_{p^*,p'^*}$ from $B^{-1}(p^*)$ to $B^{-1}(p'^*)$ such that $L(p^*) = p'^*$, and for all $(p,p') \in B^{-1}(p^*) \times B^{-1}(p'^*)$*

$$S(p, L(p)) = \|L(p) - p\| = \|p^* - p'^*\|,$$

$$S(p, L(p)) + S(L(p), p') = S(p, p').$$

Note that using matrices of example F section 4, the map L is just a homothetic transformation as illustrated below.



Proof. Note at first that $B^{-1}(p^*) = \{\tilde{p} \in \Delta(K) | \lambda(\tilde{p}) = \lambda(p^*)\}$. Since $p^*, p'^* \in P^*$, we have for all $r = 1, .., r_M$, $p^*_{|r} = p'^*_{|r} = p^*(r)$. Define the map L by

$$L(p) = \sum_{r=1}^{r_M} \lambda(p'^*)^r p_{|r}$$

It is easily seen that it defines an affine map with values in $B^{-1}(p'^*)$. Moreover,

$$\begin{aligned}
\|L(p) - p\| &= \sum_{r\,:\,\lambda(p^*)^r>0} \sum_{k\in K(r)} |(\frac{\lambda(p'^*)^r}{\lambda(p^*)^r} - 1)| p^k + \sum_{r\,:\,\lambda(p)^r=0} \lambda(p'^*)^r \sum_{k\in K(r)} p^*(r)^k \\
&= \sum_{r\,:\,\lambda(p^*)^r>0} |\lambda(p'^*)^r - \lambda(p^*)^r| + \sum_{r\,:\,\lambda(p^*)^r=0} \lambda(p'^*)^r \\
&= \sum_{r=1}^{r_M} |\lambda(p'^*)^r - \lambda(p^*)^r| = \|p^* - p'^*\| = S(p^*, p'^*).
\end{aligned}$$

For the last equality, note that by construction

$$\begin{aligned}
S(p, L(p)) &= \|\lambda(p') - \lambda(p)\|, \\
S(L(p), p') &= \sum_{r\,:\,\lambda(p')^r\lambda(p)^r>0} \lambda(p')\|p_{|r} - p'_{|r}\|,
\end{aligned}$$

so that

$$S(p, L(p)) + S(L(p), p') = S(p, p').$$

$\square$

Using the last lemma, we now prove that in some sense H is "non expansive with respect to S".

Lemma 7.13. *For all* p *and* p' *in* $\Delta(K)$ *and for all* $\mu = \sum_{n=1}^{N} \alpha_n \delta_{p_n} \in H(p)$*, there exists* $\nu = \sum_{n=1}^{N} \alpha_n \delta_{p'_n} \in H(p')$ *such that*

$$\sum_{n=1}^{N} \alpha_n S(p_n, p'_n) = S(p, p').$$

In particular $p \to H(pB)$ *is 1-Lipschitz from* $(\Delta(K), \|.\|)$ *to* $(\Delta(\Delta(K)), d_W)$*.*

Proof. Let $\mu = \sum_{n=1}^{N} \alpha_n \delta_{p_n} \in H(p)$ and $p' \in \Delta(K)$. Let us denote $p^* = pB$ and $p'^* = p'B$. Note that $p_{|r} = \sum_{n=1}^{N} \alpha_n p_{n|r}$ since for all $n = 1, .., N$, $\lambda(p_n) = \lambda(p)$. Let $L = L_{p^*,p'^*}$ be the affine map given by Lemma 7.12. Define $\tilde{p}_n = L(p_n)$ for $n = 1, .., N$, so that $L(p) = \sum_{n=1,..,N} \alpha_n \tilde{p}_n$. From Lemma 7.12, for all $p'' \in B^{-1}(p'^*)$, we have

$$S(p_n, p'') = S(p_n, \tilde{p}_n) + S(\tilde{p}_n, p'').$$

For $r = 1, .., r_M$, using Lemma 9.1 in the simplex $\Delta(K(r))$ (seen as a subset of $\Delta(K)$), there exists $(b_{r,n})_{n=1,..N} \in \Delta(K(r))$ such that $\sum_{n=1,..,N} \alpha_n b_{r,n} = p'_{|r}$ and

$$\sum_{n=1,..,N} \alpha_n \|\tilde{p}_{n|r} - b_{n,r}\| = \| \sum_{n=1,..,N} \alpha_n \tilde{p}_{n|r} - p'_{|r}\|.$$

If $\lambda(p')^r > 0$, then

$$\sum_{n=1,..,N} \alpha_n \tilde{p}_{n|r} = \sum_{n=1,..,N} \alpha_n p_{n|r} = p_{|r}.$$

Let $p'_n = \sum_{r=1,..,r_M} \lambda(p')^r b_{r,n}$, then $\sum_{n=1}^N \alpha_n p'_n = p'$ and

$$\begin{aligned}\sum_{n=1}^N \alpha_n S(\tilde{p}_n, p'_n) &= \sum_{n=1}^N \alpha_n \sum_{r\,:\,\lambda(p')^r>0} \lambda(p')^r \|\tilde{p}_{n|r} - p'_{n|r}\| \\ &= \sum_{n=1}^N \alpha_n \sum_{r\,:\,\lambda(p')^r>0} \lambda(p')^r \|\tilde{p}_{n|r} - b_{r,n}\| \\ &= \sum_{r\,:\,\lambda(p')^r>0} \lambda(p')^r \|p_{|r} - p'_{|r}\| = S(L(p), p').\end{aligned}$$

It follows that

$$\begin{aligned}\sum_{n=1}^N \alpha_n S(p_n, p'_n) &= \sum_{n=1}^N \alpha_n (S(p_n, \tilde{p}_n) + S(\tilde{p}_n, p'_n)) \\ &= S(p, L(p)) + S(L(p), p') = S(p, p').\end{aligned}$$

□

We now conclude the general proof of Proposition 5.13. Let $\varepsilon > 0$, $p' \in \Delta(K)$ and $\mu^* \in H(p)$ such that

$$\inf_{\nu \in H(q)} \left(\alpha V(\mu^*, \nu) + (1-\alpha) \int_{\Delta(K)\times\Delta(L)} f(\tilde{p}M, \tilde{q}N) d\mu^* \otimes \nu(\tilde{p}, \tilde{q}) \right) \geq \Phi_\alpha(f)(p,q) - \varepsilon$$

If $\mu^* = \sum_{n=1}^N \alpha_n \delta_{p_n}$, using Lemma 7.13, there exists $\mu' = \sum_{n=1}^N \alpha_n \delta_{p'_n} \in H(p')$ such that

$$\sum_{n=1}^N \alpha_n S(p_n, p'_n) = S(p, p').$$

Let $\nu^* \in H(q)$ such that

$$\begin{aligned}&\alpha V(\mu', \nu^*) + (1-\alpha) \int_{\Delta(K)\times\Delta(L)} f(\tilde{p}M, \tilde{q}N) d\mu' \otimes \nu^*(\tilde{p}, \tilde{q}) \\ &\leq \inf_{\nu \in H(q)} \left(\alpha V(\mu', \nu) + (1-\alpha) \int_{\Delta(K)\times\Delta(L)} f(\tilde{p}M, \tilde{q}N) d\mu' \otimes \nu(\tilde{p}, \tilde{q}) \right) + \varepsilon \leq \Phi_\alpha(f)(p', q) + \varepsilon.\end{aligned}$$

We have for all p, p', $f(pM, q) - f(p'M, q) \leq S(pM, p'M) \leq S(p, p')$, and $d_W(\mu^*, \mu') \leq$

$\sum_n \alpha_n \|p_n - p'_n\| \leq \sum_N \alpha_n S(p_n, p'_n) = S(p, p')$. Consequently, we have

$$\begin{aligned}
&\Phi_\alpha(f)(p,q) - \Phi_\alpha(f)(p',q) \\
&\qquad \leq \alpha(V(\mu^*,\nu^*) - V(\mu',\nu^*)) + (1-\alpha)\int_{\Delta(K)\times\Delta(L)} f(\tilde{p}M, \tilde{q}N) d(\mu^* - \mu') \otimes \nu^*(\tilde{p},\tilde{q}) + 2\varepsilon \\
&\qquad \leq \alpha d_W(\mu^*, \mu') + (1-\alpha)\int_{\Delta(L)} \sum_{n=1}^{N} \alpha_n (f(p_n M, \tilde{q}N) - f(p'_n M, \tilde{q}N)) d\nu^*(\tilde{q}) + 2\varepsilon \\
&\qquad \leq \alpha S(p,p') + (1-\alpha)\sum_{n=1}^{N} \alpha_n S(p_n, p'_n) + 2\varepsilon = S(p,p') + 2\varepsilon.
\end{aligned}$$

We conclude that $\Phi_\alpha(f)(p,q) - \Phi_\alpha(f)(p',q) \leq S(p,p')$. The symmetric property follows by the same method. In order to conclude the proof, it remains to prove that $\Phi_\alpha(f)$ is I-concave and II-convex on $B^{-1}(p^*) \times C^{-1}(q^*)$, but this follows directly from the fact that for all $\beta \in [0,1]$ and $p, p' \in \Delta(K)$,

$$\beta H(p) + (1-\beta)H(p') \subset H(\beta p + (1-\beta)p'),$$

and the similar property for $q, q' \in \Delta(L)$.

8 Open Questions

Note that the Lipschitz constant of 3 for the non revealing values is essentially used to obtain the uniform convergence of the non revealing value functions $\hat{v}_T$. We do not know how to prove this uniform convergence in a simpler way, neither if one can obtain a better Lipschitz constant for the non revealing values.

The Maxmin of the infinitely repeated game with incomplete information on both sides was proved in [1] to be equal to $\mathrm{cav}_I \mathrm{vex}_{II} u$. In the present model, it may be asked if the Maxmin is equal to $\mathrm{cav}_I \mathrm{vex}_{II} \hat{v}$. One may hope to prove that Player 2 can defend this quantity by combining the methods developed in the present work and the proof in [14]. It should be more complex to determine what can be guaranteed by Player 1. Even if one proves that Player 1 can guarantee the limit $\hat{v}(p,q)$ *in the non revealing game* (and consequently that the non revealing game has a uniform value), this does not imply a priori that Player 1 can guarantee $\mathrm{vex}_{II} \hat{v}$ in the original game. An other idea would be to consider the "semi-revealing" game where only Player 1 is restricted to play a non revealing strategy, then to prove that this semi-revealing game has a limit value $v'(p,q)$ that can be guaranteed by Player 1, but then there is still little hope to be able to link $\mathrm{cav}_I v'$ with $\mathrm{vex}_{II} \hat{v}$.

The results of Aumann and Maschler, and of Mertens and Zamir were extended to the case of imperfect observation of the actions. In the same way, our model can be extended to the case where instead of observing the past actions played, each player observes a stochastic signal depending only on the past actions (state independent signalling). The notion of non-revealing strategies still makes sense, so conceptually one may hope to use the same ideas to prove the convergence of the value functions in this case. Note that a difficulty in this model is that even if we fix the strategy of a player, say Player 1, this player is not able to compute along the play the belief of Player 2 about the current state in K. And when introducing signals, it is often the case that one has to construct strategies by blocks sufficiently long to allow for statistical tests. How these blocks would interfer with the blocks where non revealing strategies are used as in point (5) of the proof of Theorem 6.1 is unclear, and this seems a highly technical problem.

An important extension of repeated games with incomplete information on both sides is the case of general, possibly correlated, probabilities for the initial pair of states (k, l). The results of Mertens and Zamir [14] were actually stated within this more general framework. This suggests the following generalization of our model: we are given an initial probability $\pi \in \Delta(K \times L)$ and a Markov chain $z_t = (k_t, l_t)$ with values in $K \times L$. At each stage, Player 1 observes k_t and Player 2 observes l_t. It is still possible to define the appropriate notion of convex and concave hull as done by Mertens and Zamir. However, the definition a non-revealing strategy should be adapted. Indeed, even if Player 1 does not use his information on the state, the beliefs of Player 2 on the recurrence classes of the chain z_t may evolve. This is left for future research.

9 Appendix

Lemma 9.1. *(Laraki [11]) K being a finite set, consider the simplex $\Delta(K) = \{(p^k)_{k \in K} \in I\!R_+^K, \sum_{k \in K} p^k = 1\}$ endowed with the L^1-norm $\|p - p'\| = \sum_{k \in K} |p^k - p'^k|$.*

Consider a convex combination $p = \sum_{s \in S} \lambda_s p_s$ in $\Delta(K)$, with S finite, $\lambda_s \geq 0$ and $\sum_{s \in S} \lambda_s = 1$. Then for every p' in $\Delta(K)$ there exists $(p'_s)_{s \in S}$ in $\Delta(K)^S$ such that:

$$p' = \sum_{s \in S} \lambda_s p'_s \text{ and } \sum_{s \in S} \lambda_s \|p_s - p'_s\| = \|p - p'\|.$$

Proof[2]: For each k in K, denote by e_k the unit vector $(e_k^{k'})_{k'}$ in $\Delta(K)$ such that $e_k^k = 1$. Define $K^+ = \{k \in K, p'^k \geq p^k\}$ and $K^- = \{k \in K, p'^k < p^k\}$. Define also: $A =_{def} \sum_{k \in K^+} (p'^k - p^k) = \sum_{k \in K^-} (p^k - p'^k) = \frac{1}{2}\|p - p'\|$, and introduce $\theta \in \Delta(K^+)$ by $\theta^k = (p'^k - p^k)/A$ for all k in K^+. One can now define for all s:

$$p'_s = p_s + \sum_{k \in K^-} \frac{p_s^k}{p^k} \left(p^k - p'^k\right) (\theta - e_k).$$

1) For all k in K^-, we have $p'^k_s = p_s^k p'^k / p^k \geq 0$, and $p'_s \in \Delta(K)$.
2) $\sum_s \lambda_s p'_s = p + \sum_{k \in K^-} (p^k - p'^k)(\theta - e_k) = p + A\theta + \sum_{k \in K^-} (-p^k + p'^k) e_k = p + p' - p = p'$.
3) $p'_s - p_s = \theta \left(\sum_{k \in K^-} \frac{p_s^k}{p^k} (p^k - p'^k) \right) - \sum_{k \in K^-} \frac{p_s^k}{p^k} (p^k - p'^k) e_k$. We have a difference of vectors with disjoint support, so the norms add up and: $\|p_s - p'_s\| = 2 \sum_{k \in K^-} \frac{p_s^k}{p^k} (p^k - p'^k)$. This is true for each s, and we obtain:

$$\sum_{s \in S} \lambda_s \|p_s - p'_s\| = 2 \sum_{k \in K^-} \left(p^k - p'^k\right) = \|p - p'\|. \qquad \square$$

---

[2] The proof can be deduced from Lemma 8.2 in [11], which deals with general measurable spaces. The simpler but less general proof given here is easily deduced from the version written in R. Laraki's PhD thesis.

References

[1] R. J. Aumann and M. Maschler, *Repeated games with incomplete information*, with the collaboration of R. Stearns, Cambridge, MA: MIT Press, 1995.

[2] D. Blackwell, *Equivalent comparisons of experiments*, The Annals of Mathematical Statistics,1953, **24**, 265-272.

[3] P. Cardaliaguet, R. Laraki and S. Sorin, *A Continuous Time Approach for the Asymptotic Value in Two-Person Zero-Sum Repeated Games*, SIAM Journal on Control and Optimization, 2012, **50**, 1573-1596.

[4] B. DE Meyer and D. Rosenberg, *"Cav u" and the Dual Game*, Mathematics of Operations Research, 1999, **24**, 619-626.

[5] F. Gensbittel, *Extensions of the Cav(u) theorem for repeated games with one-sided information*, 2012, Preprint, Hal-00745575

[6] P. Gordon,*Theory of finite Markov chains*, in French, Dunod, Paris, 1964.

[7] S. Hart, *Nonzero-sum two-person repeated games with incomplete information*, Mathematics of Operations Research, 1985, **10**, 117-153

[8] M. Heuer, *Asymptotically optimal strategies in repeated games with incomplete information*, International Journal of Game Theory, 1992, **20**, 377-392.

[9] J. Horner, D. Rosenberg, E. Solan, and N. Vieille, *On a Markov Games with One-Sided Information*, Operations Research,2010, **58**, 1107-1115.

[10] E. Kohlberg, *Optimal strategies in repeated games with incomplete information*, International Journal of Game Theory, 1975, **4**, 7-24.

[11] R. Laraki, *On the Regularity of the Convexification Operator on a Compact Set*, Journal of Convex Analysis, 2004, **11**, 209-234.

[12] R. Laraki, *Repeated games with lack of information on one side: the dual differential approach*, Mathematics of Operations Research, 2002, **27**, 419-440.

[13] R. Laraki, *The splitting game and applications*, International Journal of Game Theory, 2002, **30**, 359-376.

[14] J. F. Mertens and S. Zamir, *The value of two-person zero-sum repeated games with lack of information on both sides*, International Journal of Game Theory,1971-1972, **1**, 39-64.

[15] J. F. Mertens and S. Zamir, *A duality theorem on a pair of simultaneous functional equations*, Journal of Mathematical Analysis and Applications, 1977, **60**, 550-558.

[16] A. Neyman, *Existence of optimal strategies in Markov games with incomplete information*, International Journal of Game Theory, 2008, **37**, 581-596.

[17] J. R. Norris, *Markov chains*, Cambridge Series in Statistical and Probabilistic Mathematics, Cambridge University Press, 1997.

[18] J. Renault, *2-player repeated games with lack of information on one side and state independent signalling*, Mathematics of Operations Research, 2000, **25**, 552-572.

[19] J. Renault, *3-player repeated games with lack of information on one side*, International Journal of Game Theory, 2001, **30**, 221-246.

[20] J. Renault, *The value of Markov chain games with lack of information on one side*, Mathematics of Operations Research, 2006, **31**, 490-512.

[21] J. Renault, *The value of repeated games with an informed controller*, Mathematics of Operations Research, 2012, **37**, 154-179.

[22] D. Rosenberg and S. Sorin, *An operator approach to zero-sum repeated games*, Israel Journal of Mathematics, 2001, **121**, 221-246.

[23] R.S. Simon, S. and Spież, and H. Toruńczyk, *The existence of equilibria in certain games, separation for families of convex functions and a theorem of Borsuk-Ulam type*, Israel Journal of Mathematics, 1995, **92**, 1-21.

[24] M. Sion, *On general minimax theorems*, Pacific Journal of Mathematics, 1995, **8**, 171-176.

[25] S. Sorin, *A first course on zero-sum repeated games*, Mathématiques & Applications, Springer, 2002.

[26] S.Sorin, *Some results on the existence of Nash equilibria for non-zero sum games with incomplete information*, International Journal of Game Theory, 1983, **12**, 193-205.

[27] S.Sorin, *On a pair of simultaneous functional equations*, Journal of mathematical analysis and applications, 1984, **98**, 296-303.

[28] C. Villani, *Topics in Optimal Transportation.* Graduate Studies in Mathematics, Vol. 58. American Mathematical Society, 2003.

[29] S. Zamir, *Repeated Games of Incomplete Information: Zero-Sum*, Handbook of Game Theory with Economic Applications, edited by Aumann and Hart. Vol I, chapter 6, 1992.